

ADVISORY COMMITTEE ON THE MICROBIOLOGICAL SAFETY OF FOOD

Items of possible interest from the literature

A list of items from the literature which may be of interest to members is attached.

**Secretariat
March 2007**

Bacillus

Fricker M, Messelhauser U, Busch U, Scherer S, Ehling-Schultz M. Diagnostic real-time PCR assays for the detection of emetic *Bacillus cereus* strains in foods and recent foodborne outbreaks. *Appl Environ Microbiol* 2007 Jan 26. Epub ahead of print.

Park DK, Bitton G, Melker R. Microbial inactivation by microwave radiation in the home environment. *J Environ Health* 2006; **69(5)**: 17-24; quiz 39-40.

Vissers MM, Te Giffel MC, Driehuis F, De Jong P, Lankveld JM. Predictive modeling of *Bacillus cereus* spores in farm tank milk during grazing and housing periods. *J Dairy Sci* 2007; **90(1)**: 281-92.

Rasko DA, Rosovitz MJ, Okstad OA, Fouts DE, Jiang L, Cer RZ, Kolsto AB, Gill SR, Ravel J. Complete sequence analysis of novel plasmids from emetic and periodontal *Bacillus cereus* isolates reveals a common evolutionary history among the *B. cereus*-group plasmids, including *Bacillus anthracis* pXO1. *J Bacteriol* 2007; **189(1)**: 52-64.

Van der Auwera GA, Timmery S, Hoton F, Mahillon J. Plasmid exchanges among members of the *Bacillus cereus* group in foodstuffs. *Int J Food Microbiol* 2007; **113(2)**: 164-72.

From C, Homazabal V, Granum PE. Food poisoning associated with pubilacidin-producing *Bacillus pumilis* in rice. *Int J Food Microbiol* 2007 Jan 13. Epub ahead of print.

Campylobacter

Richardson G, Thomas DR, Smith RM, Nehaul L, Ribeiro CD, Brown AG, Salmon RL. A community outbreak of *Campylobacter jejuni* infection from a chlorinated public water supply. *Epidemiol Infect* 2007 Feb 9: 1-8. Epub ahead of print.

Ruiz-Palacios GM. The health burden of campylobacter infection and the impact of antimicrobial resistance: playing chicken. *Clin Infect Dis* 2007; **44(5)**: 701-3.

Tsuei AC, Carey-Smith GV, Hudson JA, Billington C, Heinemann JA. Prevalence and numbers of coliphages and *Campylobacter jejuni* bacteriophages in New Zealand foods. *Int J Food Microbiol* 2007 Jan 16. Epub ahead of print.

Singer RS, Cox LA Jr, Dickson JS, Hurd HS, Phillips I, Miller GY. Modeling the relationship between food animal health and human foodborne illness. *Prev Vet Med* 2007 Jan 29. Epub ahead of print.

- Hanel CM, Atanassova V. Impact of different storage factors on the survivability of *Campylobacter jejuni* in turkey meat. FEMS Immunol Med Microbiol 2007; **49(1)**: 146-8.
- Atanassova V, Reich F, Beckmann L, Klein G. Prevalence of *Campylobacter* spp. in turkey meat from a slaughterhouse and in turkey meat retail products. FEMS Immunol Med Microbiol 2007; **49(1)**: 141-5.
- Corcoran AT, Moran AP. Influence of growth conditions on diverse polysaccharide production by *Campylobacter jejuni*. FEMS Immunol Med Microbiol 2007; **49(1)**: 124-32.
- Rasschaert G, Houf K, De Zutter L. External contamination of *Campylobacter*-free flocks after transport in cleaned and disinfected containers. J Food Prot 2007; **70(1)**: 40-6.
- Luber P, Bartelt E. Enumeration of *Campylobacter* spp. on the surface and within chicken breast fillets. J Appl Microbiol 2007; **102(2)**: 313-8.
- Boysen L, Knochel S, Rosenquist H. Survival of *Campylobacter jejuni* in different gas mixtures. FEMS Microbiol Lett 2007; **266(2)**:152-7.
- de Zoete MR, van Putten JP, Wagenaar JA. Vaccination of chickens against *Campylobacter*. Vaccine 2006 Dec 15. Epub ahead of print.
- Kuwabara S. Guillain-barre syndrome. Curr Neurol Neurosci Rep 2007; **7(1)**: 57-62.
- Johannessen GS, Johnsen G, Okland M, Cudjoe KS, Hofshagen M. Enumeration of thermotolerant *Campylobacter* spp. from poultry carcasses at the end of the slaughter-line. Lett Appl Microbiol 2007; **44(1)**: 92-7.
- Fullerton KE, Ingram LA, Jones TF, Anderson BJ, McCarthy PV, Hurd S, Shiferaw B, Vugia D, Haubert N, Hayes T, Wedel S, Scallan E, Henao O, Angulo FJ. Sporadic *campylobacter* infection in infants: a population-based surveillance case-control study. Pediatr Infect Dis J 2007; **26(1)**: 19-24.
- Lindmark H, Diedrich IC, Andersson L, Lindqvist R, Engvall EO. Distribution of *Campylobacter* genotypes on broilers during slaughter. J Food Prot 2006; **69(12)**: 2902-7.
- Chen L, Geys H, Cawthraw S, Havelaar A, Teunis P. Dose response for infectivity of several strains of *Campylobacter jejuni* in chickens. Risk Anal 2006; **26(6)**: 1613-21.
- Byrd J, Bailey RH, Wills R, Nisbet D. Recovery of *Campylobacter* from commercial hatchery trayliners. Poult Sci 2007; **86(1)**: 26-9.
- James C, James SJ, Hannay N, Purnell G, Barbedo-Pinto C, Yaman H, Araujo M, Gonzalez ML, Calvo J, Howell M, Corry JE. Decontamination of

poultry carcasses using steam or hot water in combination with rapid cooling, chilling or freezing of carcass surfaces. *Int J Food Microbiol* 2006; Nov 29. Epub ahead of print.

Stafford RJ, Schluter P, Kirk M, Wilson A, Unicomb L, Ashbolt R, Gregory J; the OzFoodNet Working Group. A multi-centre prospective case-control study of campylobacter infection in persons aged 5 years and older in Australia. *Epidemiol Infect* 2006 Nov 30; 1-11. Epub ahead of print.

Houf K, Stephan R. Isolation and characterization of the emerging foodborne pathogen *Arcobacter* from human stool. *J Microbiol Methods* 2007; **68(2)**: 408-13.

Karenlampi R, Rautelin H, Schonberg-Norio D, Paulin L, Hanninen ML. Longitudinal study of Finnish *Campylobacter jejuni* and *C. coli* isolates from humans, using multilocus sequence typing, including comparison with epidemiological data and isolates from poultry and cattle. *Appl Environ Microbiol* 2007; **73(1)**: 148-55.

Duffy L, Dykes GA. Growth temperature of four *Campylobacter jejuni* strains influences their subsequent survival in food and water. *Lett Appl Microbiol* 2006; **43(6)**: 596-601.

Whyte R, Hudson JA, Graham C. *Campylobacter* in chicken livers and their destruction by pan frying. *Lett Appl Microbiol* 2006; **43(6)**: 591-5.

Allen VM, Bull SA, Corry JE, Domingue G, Jorgensen F, Frost JA, Whyte R, Gonzalez A, Elviss N, Humphrey TJ. *Campylobacter* spp. contamination of chicken carcasses during processing in relation to flock colonisation. *Int J Food Microbiol* 2007; **113(1)**: 54-61.

Son I, Englen MD, Berrang ME, Fedorka-Cray PJ, Harrison MA. Prevalence of *Arcobacter* and *Campylobacter* on broiler carcasses during processing. *Int J Food Microbiol* 2007; **113(1)**: 16-22.

Smith A, Reacher M, Smerdon W, Adak GK, Nichols G, Chalmers RM. Outbreaks of waterborne infectious intestinal disease in England and Wales 1992-2003. *Epidemiol Infect* 2006; **134(6)**: 1141-9.

van de Giessen AW, Bouwknecht M, Dan-Deisz WD, van Pelt W, Wannet WJ, Visser G. Surveillance of *Salmonella* spp. and *Campylobacter* spp. in poultry production flocks in The Netherlands. *Epidemiol Infect* 2006; **134(6)**: 1266-75.

McGill K, Cowley D, Moran L, Scates P, O'Leary A, Madden RH, Carroll C, McNamara E, Moore JE, Fanning S, Collins JD, Whyte P. Antibiotic resistance of retail food and human *Campylobacter* isolates on the island of Ireland from 2001-2002. *Epidemiol Infect* 2006; **134(6)**: 1282-91.

Clostridium botulinum

Webb MD, Pin C, Peck MW, Stringer SC. Historical and contemporary NaCl concentrations affect the duration and distribution of lag times from individual spores of non-proteolytic *Clostridium botulinum*. *Appl Environ Microbiol* 2007 Feb 2. Epub ahead of print.

Centers for Disease Control and Prevention (CDC). Foodborne botulism from home-prepared fermented tofu-California, 2006. *MMWR Morb Mortal Wkly Rep* 2007; **56(5)**: 96-7.

Heffron A, Poxton IR. A PCR approach to determine the distribution of toxin genes in closely related *Clostridium* species: *Clostridium botulinum* type C and D neurotoxins and C2 toxin, and *Clostridium novyi* {alpha} toxin. *J Med Microbiol* 2007; **56(Pt 2)**: 196-201.

Hill KK, Smith TJ, Helma CH, Ticknor LO, Foley BT, Svensson RT, Brown JL, Johnson EA, Smith LA, Okinaka RT, Jackson PJ, Marks JD. Genetic diversity among *Botulinum* Neurotoxin-producing clostridial strains. *J Bacteriol* 2007; **189(3)**: 818-32.

Dolly JO, Aoki KR. The structure and mode of action of different botulinum toxins. *Eur J Neurol* 2006; **13 Suppl 4**: 1-9.

McLauchlin J, Grant KA, Little CL. Food-borne botulism in the United Kingdom. *J Public Health (Oxf.)* 2006; **28(4)**: 337-42.

Cryptosporidium

Abubakar I, Aliyu Sh, Arumugam C, Hunter P, Usman N. Prevention and treatment of cryptosporidiosis in immunocompromised patients. *Cochrane Database Syst Rev* 2007; **1**: CD004932.

Thompson HP, Dooley JS, Kenny J, McCoy M, Lowery CJ, Moore JE, Xiao L. Genotypes and subtypes of *Cryptosporidium* spp. in neonatal calves in Northern Ireland. *Parasitol Res* 2007; **100(3)**: 619-24.

Schets FM, van den Berg HH, Engels GB, Lodder WJ, de Roda Husman AM. *Cryptosporidium* and *Giardia* in commercial and non-commercial oysters (*Crassostrea gigas*) and water from the Oosterschelde, The Netherlands. *Int J Food Microbiol* 2007; **113(2)**: 189-94.

Graczyk TK, Lewis EJ, Glass G, Dasilva AJ, Tamang L, Girouard AS, Curriero FC. Quantitative assessment of viable *Cryptosporidium parvum* load in commercial oysters (*Crassostrea virginica*) in the Chesapeake Bay. *Parasitol Res* 2007; **100(2)**: 247-53.

***E. coli* O157**

Franz E, Klerks MM, De Vos OJ, Termorshuizen AJ, van Bruggen AH. Farm prevalence of STEC stx1, stx2, eaeA and rfbE genes and survival of *E. coli* O157: H7 in manure from organic and low-input conventional dairy farms. *Appl Environ Microbiol* 2007 Feb 2. Epub ahead of print.

Azizoglu RO, Drake M. Impact of antibiotic stress on acid and heat tolerance and virulence factor expression of *Escherichia coli* O157: H7. *J Food Prot* 2007; **70(1)**: 194-9.

Schaffner DW, Schaffner KM. Management of risk of microbial cross-contamination from uncooked frozen hamburgers by alcohol-based hand sanitizer. *J Food Prot* 2007; **70(1)**: 109-13.

Doane CA, Pangloli P, Richards HA, Mount JR, Golden DA, Draughon FA. Occurrence of *Escherichia coli* O157:H7 in diverse farm environments. *J Food Prot* 2007; **70(1)**: 6-10.

Bergholz TM, Whittam TS. Variation in acid resistance among enterohaemorrhagic *Escherichia coli* in a simulated gastric environment. *J Appl Microbiol* 2007; **102(2)**: 352-62.

Kresse AU, Rienacker I, Valle AM, Steinruck H, Claus H, Payne SM, Tschape H, Williams PH, Reissbrodt R. Enterohaemorrhagic *Escherichia coli* O157 and non-O157 serovars differ in their mechanisms for iron supply. *Int J Med Microbiol* 2007; **297(1)**: 9-15.

Stacey KF, Parsons DJ, Christiansen KH, Burton CH. Assessing the effect of interventions on the risk of cattle and sheep carrying *Escherichia coli* O157:H7 to the abattoir using a stochastic model. *Prev Vet Med* 2007 Jan 2. Epub ahead of print.

Manago S, Kishikawa K, Tokunaga H, Funatsumaru S, Kubo Y, Sonoda M, Yoshihara T, Nasu F, Kasahara K. Outbreak of enterohemorrhagic *Escherichia coli* O157 attributed to a grilled-meat restaurant. *Jpn J Infect Dis* 2006; **59(6)**: 407-8.

Alam MJ, Zurek L. Seasonal prevalence of *Escherichia coli* O157:H7 in beef cattle feces. *J Food Prot* 2006; **69(12)**: 3018-20.

Islam MA, Heuvelink AE, Talukder KA, Zwietering MH, de Boer E. Evaluation of immunomagnetic separation and PCR for the detection of *Escherichia coli* O157 in animal feces and meats. *J Food Prot* 2006; **69(12)**: 2865-9.

Echeverry A, Loneragan GH, Brashears MM. Survival of *Escherichia coli* O157:H7 in bovine feces over time under various temperature conditions. *J Food Prot* 2006; **69(12)**: 2851-5.

Woerner DR, Ransom JR, Sofos JN, Dewell GA, Smith GC, Salman MD, Belk KE. Determining the prevalence of *Escherichia coli* O157 in cattle and beef from the feedlot to the cooler. *J Food Prot* 2006; **69(12)**: 2824-7.

Yang HH, Morrow JB, Grasso D, Vinopal RT, Smets BF. Intestinal versus external growth conditions change the surficial properties in a collection of environmental *Escherichia coli* isolates. *Environ Sci Technol* 2006; **40(22)**: 6976-82.

Voetsch AC, Kennedy MH, Keene WE, Smith KE, Rabatsky-Ehr T, Zansky S, Thomas SM, Mohle-Boetani J, Sparling PH, McGavern MB, Mead PS. Risk factors for sporadic Shiga toxin-producing *Escherichia coli* O157 infections in FoodNet sites, 1999-2000. *Epidemiol Infect* 2006 Dec 6; 1-8. Epub ahead of print.

Lejeune JT, Wetzel AN. Preharvest control of *Escherichia coli* O157 in cattle. *J Anim Sci* 2006 Dec 4. Epub ahead of print.

Besser TE, Shaikh N, Holt NJ, Tarr PI, Konkel ME, Malik-Kale P, Walsh CW, Whittam TS, Bono JL. Greater diversity of shiga toxin-encoding bacteriophage insertion sites among *Escherichia coli* O157:H7 isolates from cattle than in those from humans. *Appl Environ Microbiol* 2007; **73(3)**: 671-9.

Halliday JE, Chase-Topping ME, Pearce MC, McKendrick IJ, Allison L, Fenlon D, Low C, Mellor DJ, Gunn GJ, Woolhouse ME. Herd-level risk factors associated with the presence of phage type 21/28 *E. coli* O157 on Scottish cattle farms. *BMC Microbiol* 2006; **6**: 99.

[no authors listed] *E. coli* O157:H7 outbreak in the United States associated with bagged fresh spinach. *Can Commun Dis Rep* 2006; **32(22)**: 272.

Wood JC, McKendrick IJ, Gettinby G. A simulation model to assess herd-level intervention strategies against *E. coli* O157. *Epidemiol Infect* 2006 Nov 17: 1-16. Epub ahead of print.

Johnson KE, Thorpe CM, Sears CL. The emerging clinical importance of non-O157 shiga toxin-producing *Escherichia coli*. *Clin Infect Dis* 2006; **43(12)**: 1587-95.

Hoffman MA, Menge C, Casey TA, Laegreid W, Bosworth BT, Dean-Nystrom EA. Bovine immune response to shiga-toxigenic *Escherichia coli* O157: H7. *Clin Vaccine Immunol* 2006; **13(12)**: 1322-7.

Ahn J, Grun IU, Mustapha A. Effects of plant extracts on microbial growth, color change, and lipid oxidation in cooked beef. *Food Microbiol* 2007; **24(1)**: 7-14.

Hepatitis A

Shieh YC, Khudyakov YE, Xia G, Ganova-Raeva LM, Khambaty FM, Woods JW, Veazey JE, Motes ML, Glatzer MB, Bialek SR, Fiore AE. Molecular confirmation of oysters as the vector for hepatitis A in a 2005 multistate outbreak. *J Food Prot* 2007; **70(1)**: 145-50.

Croci L, Losio MN, Suffredini E, Pavoni E, Di Pasquale S, Fallacara F, Arcangeli G. Assessment of human enteric viruses in shellfish from the northern Adriatic sea. *Int J Food Microbiol* 2006 Dec 27. Epub ahead of print.

Hepatitis E

Seminati C, Mateu E, Peralta B, de Deus N, Martin M. Distribution of hepatitis E virus infection and its prevalence in pigs on commercial farms in Spain. *Vet J* 2007 Feb 3. Epub ahead of print.

Herremans M, Vennema H, Bakker J, van der Veer B, Duizer E, Benne CA, Waar K, Hendrixks B, Schneeberger P, Blaauw G, Kooiman M, Koopmans MP. Swine-like hepatitis E viruses are a cause of unexplained hepatitis in the Netherlands. *J Viral Hepat* 2007; **14(2)**: 140-6.

de Deus N, Seminati C, Pina S, Mateur E, Martin M, Segales J. Detection of hepatitis E virus in liver, mesenteric lymph node, serum, bile and faeces of naturally infected pigs affected by different pathological conditions. *Vet Microbiol* 2007; **119(2-4)**: 105-14.

Listeria monocytogenes

Glass K, Preston D, Veessenmeyer J. Inhibition of *Listeria monocytogenes* in turkey and pork-beef bologna by combinations of sorbate, benzoate, and propionate. *J Food Prot* 2007; **70(1)**: 214-7.

Mejlholm O, Dalgaard P. Modeling and predicting the growth boundary of *Listeria monocytogenes* in lightly preserved seafood. *J Food Prot* 2007; **70(1)**: 70-84.

Varma JK, Samuel MC, Marcus R, Hoekstra RM, Medus C, Segler S, Anderson BJ, Jones TF, Shiferaw B, Haubert N, Megginson M, McCarthy PV, Graves L, Gilder TV, Angulo FJ. *Listeria monocytogenes* infection from foods prepared in a commercial establishment: a case-control study of potential sources of sporadic illness in the United States. *Clin Infect Dis* 2007; **44(4)**: 521-8.

Voetsch AC, Angulo FJ, Jones TF, Moore MR, Nadon C, McCarthy P, Shiferaw B, Megginson MB, Hurd S, Anderson BJ, Cronquist A, Vugia DJ, Medus C, Segler S, Graves LM, Hoekstra RM, Griffin PM; Centers for Disease Control and Prevention Emerging Infections Program Foodborne Diseases Active Surveillance Network Working Group. Reduction in the incidence of

invasive listeriosis in foodborne diseases active surveillance network sites, 1996-2003. Clin Infect Dis 2007; **44(4)**: 513-20.

Hong E, Doumith M, Duperrier S, Giovannacci I, Morvan A, Glaser P, Buchrieser C, Jacquet C, Martin P. Genetic diversity of *Listeria monocytogenes* recovered from infected persons and pork, seafood and dairy products on retail sale in France during 2000 and 2001. Int J Food Microbiol 2006 Dec 22. Epub ahead of print.

Vorst KL, Todd EC, Ryser ET. Transfer of *Listeria monocytogenes* during slicing of turkey breast, bologna, and salami with simulated kitchen knives. J Food Prot 2006; **69(12)**: 2939-46.

Vermeulen A, Gysemans KP, Bernaerts K, Geeraerd AH, Van Impe JF, Debevere J, Devlieghere F. Influence of pH, water activity and acetic acid concentration on *Listeria monocytogenes* at 7 degrees C: Data collection for the development of a growth/no growth model. Int J Food Microbiol 2006 Dec 19. Epub ahead of print.

Jallewar PK, Kalorey DR, Kurkure NV, Pande VV, Barbuddhe SB. Genotypic characterization of *Listeria* spp. isolated from fresh water fish. Int J Food Microbiol 2007; **114(1)**: 120-3.

Carrasco E, Valero A, Perez-Rodriguez F, Garcia-Gimeno RM, Zurera G. Management of microbiological safety of ready-to-eat meat products by mathematical modelling: *Listeria monocytogenes* as an example. Int J Food Microbiol 2006 Nov 29. Epub ahead of print.

Tham W, Valladares GL, Helmersson S, Osterlund A, Danielsson-Tham ML. More than one variant of *Listeria monocytogenes* isolated from each of two human cases of invasive listeriosis. Epidemiol Infect 2006 Nov 17: 1-3. Epub ahead of print.

Crepet A, Albert I, Dervin C, Carlin F. Estimation of microbial contamination of food from prevalence and concentration data: application of *Listeria monocytogenes* in fresh vegetables. Appl Environ Microbiol 2007; **73(1)**: 250-8.

Brouwer MC, van de Beek D, Heckenberg SG, Spanjaard L, de Gans J. Community-acquired *Listeria monocytogenes* meningitis in adults. Clin Infect Dis 2006; **43(10)**: 1233-8.

Gandhi M, Chikindas ML. *Listeria*: A foodborne pathogen that knows how to survive. Int J Food Microbiol 2007; **113(1)**: 1-15.

Norovirus

Fell G, Boyens M, Baumgarte S. Frozen berries as a risk factor for outbreaks of norovirus gastroenteritis. Results of an outbreak investigation in the

summer of 2005 in Hamburg. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2007 Jan 24. Epub ahead of print.

Wang QH, Costantini V, Saif LJ. Porcine enteric caliciviruses: Genetic and antigenic relatedness to human caliciviruses, diagnosis and epidemiology. Vaccine 2006 Dec 29. Epub ahead of print.

Cheetham S, Souza M, McGregor R, Meulia T, Wang Q, Saif LJ. Binding patterns of human norovirus-like particles to buccal and intestinal tissues of gnotobiotic pigs in relation to A/H histo-blood group antigen expression. J Virol 2007 Jan 10. Epub ahead of print.

Croci L, Losio MN, Suffredini E, Pavoni E, Di Pasquale S, Fallacara F, Arcangeli G. Assessment of human enteric viruses in shellfish from the northern Adriatic sea. Int J Food Microbiol 2006 Dec 27. Epub ahead of print.

Murata T, Katsushima N, Mizuta K, Muraki Y, Hongo S, Matsuzaki Y. Prolonged norovirus shedding in infants ≤ 6 months of age with gastroenteritis. Pediatr Infect Dis J 2006; **26(1)**: 46-9.

Schultz AC, Saadbye P, Hoorfar J, Norrung B. Comparison of methods for detection of norovirus in oysters. Int J Food Microbiol 2006 Dec 18. Epub ahead of print.

Sano D, Ueki Y, Watanabe T, Omura T. Genetic variation in the conservative gene region of Norovirus genogroup II strains in environmental and stool samples. Environ Sci Technol 2006; **40(23)**: 7423-7.

Kingsley DH, Holliman DR, Calci KR, Chen H, Flick GJ. Inactivation of a norovirus by high-pressure processing. Appl Environ Microbiol 2007; **73(2)**: 581-5.

Shirato-Horikoshi H, Ogawa S, Wakita T, Takeda N, Hansman GS. Binding activity of norovirus and sapovirus to histo-blood group antigens. Arch Virol 2006 Nov 27. Epub ahead of print.

Butot S, Putallaz T, Sanchez G. Procedure for rapid concentration and detection of enteric viruses from berries and vegetables. Appl Environ Microbiol 2007; **73(1)**: 186-92.

Tian P, Bates AH, Jensen HM, Mandrell RE. Norovirus binds to blood group A-like antigens in oyster gastrointestinal cells. Lett Appl Microbiol 2006; **43(6)**: 645-51.

Salmonella

de Abreu Correa A, Albarnaz JD, Moresco V, Poli CR, Teixeira AL, Oliveira Simoes CM, Monte Barardi CR. Depuration dynamics of oysters (*Crassostrea gigas*) artificially contaminated by *Salmonella enterica* serovar Typhimurium. Mar Environ Res 2006 Dec 21. Epub ahead of print.

Gupta SK, Nalluswami K, Snider C, Perch M, Balasegaram M, Burmeister D, Lockett J, Sandt C, Hoekstra RM. Outbreak of *Salmonella Braenderup* infections associated with Roma tomatoes, northeastern United States, 2004: a useful method for subtyping exposures in field investigations. *Epidemiol Infect* 2007 Feb 5: 1-9. Epub ahead of print.

Avila LA, Nascimento VP, Salle CT, Moraes HL. Effects of probiotics and maternal vaccination on *Salmonella enteritidis* infection in broiler chicks. *Avian Dis* 2006; **50(4)**: 608-12.

Dominguez A, Torner N, Ruiz L, Martinez A, Bartolome R, Sulleiro E, Teixido A, Plasencia A. Foodborne Salmonella-caused outbreaks in Catalonia (Spain), 1990 to 2003. *J Food Prot* 2007; **70(1)**: 209-13.

Iturriaga MH, Tamplin ML, Escartin EF. Colonization of tomatoes by *Salmonella montevideo* is affected by relative humidity and storage temperature. *J Food Prot* 2007; **70(1)**: 30-4.

Adriansen C, De Greve H, Tian JQ, De Craeye S, Gubbels E, Eeckhaut V, Van Immerseel F, Ducatelle R, Kumar M, Hernalsteens JP. Live *Salmonella enterica* serovar *Enteritidis* vaccine allowing serological differentiation between vaccinated and infected animals. *Infect Immun* 2007 Jan 29. Epub ahead of print.

Brouard C, Espie E, Weill FX, Kerouanton A, Brisavois A, Forgue AM, Vaillant V, de Valk H. Two consecutive large outbreaks of *Salmonella enterica* serotype *Agona* infections in infants linked to the consumption of powdered infant formula. *Pediatr Infect Dis J* 2007; **26(2)**: 148-152.

Madden RH, Murray KA, Gilmour A. Carriage of four bacterial pathogens by beef cattle in Northern Ireland at time of slaughter. *Lett Appl Microbiol* 2007; **44(2)**: 115-9.

Santiesteban-Lopez A, Palou E, Lopez-Malo A. Susceptibility of food-borne bacteria to binary combinations of antimicrobials at selected a and pH. *J Appl Microbiol* 2007; **102(2)**: 486-97.

Asai T, Ishihara K, Harada K, Kojima A, Tamura Y, Sato S, Takahashi T. Long-term prevalence of antimicrobial-resistant *Salmonella enterica* Subspecies *enterica* Serovar *Infantis* in the broiler chicken industry in Japan. *Microbiol Immunol* 2007; **51(1)**: 111-5.

Little CL, Walsh S, Hucklesby L, Surman-Lee S, Pathak K, Hall Y, de Pinna E, Threlfall EJ, Maund A, Chan CH. *Salmonella* contamination in non-UK produced shell eggs on retail sale in some regions of England. *Euro Surveill* 2006; **11(11)**: E061123.4.

Freeman AF, Holland SM. Persistent bacterial infections and primary immune disorders. *Curr Opin Microbiol* 2007 Jan 5. Epub ahead of print.

Noel H, Dominguez M, Weill FX, Brisabois A, Duchazeaubeneix C, Kerouanton A, Delmas G, Pihier N, Couturier E. Outbreak of *Salmonella enterica* serotype Manhattan infection associated with meat products, France, 2005. *Euro Surveill* 2006; **11(11)**. Epub ahead of print.

Davison HC, Sayers AR, Smith RP, Pascoe SJ, Davies RH, Weaver JP, Evans SJ. Risk factors associated with the *salmonella* status of dairy farms in England and Wales. *Vet Rec* 2006; **159(26)**: 871-80.

Boyle EC, Bishop JL, Grassl GA, Finlay BB. *Salmonella*: From pathogenesis to therapeutics. *J Bacteriol* 2006 Dec 22. Epub ahead of print.

Chiewchan N, Pakdee W, Devahastin S. Effect of water activity on thermal resistance of *Salmonella krefeld* in liquid medium and on rawhide surface. *Int J Food Microbiol* 2007; **114(1)**: 43-9.

Ebuchi S, Baba A, Uryu K, Hiwaki H. Two outbreaks caused by *Salmonella derby* and *S. Anatum* at grilled-meat restaurants in fukuoka city. *Jpn J Infect Dis* 2006; **59(6)**: 405-6.

Lee SY, Chung HJ, Shin JH, Dougherty RH, Kangi DH. Survival and growth of foodborne pathogens during cooking and storage of oriental-style rice cakes. *J Food Prot* 2006; **69(12)**: 3037-42.

Murase T, Chiba K, Sato T, Otsuki K, Holt PS. Effects of different molting procedures on incidence of *Salmonella* infection in flocks of naturally contaminated laying hens in a commercial egg-producing farm by detection of yolk antibodies to *Salmonella* in eggs. *J Food Prot* 2006; **69(12)**: 2883-8.

Lindblad M, Lindmark H, Lambertz ST, Lindqvist R. Microbiological baseline study of broiler chickens at Swedish slaughterhouses. *J Food Prot* 2006; **69(12)**: 2875-82.

McMeechan A, Roberts M, Cogan TA, Jorgensen F, Stevenson A, Lewis C, Rowley G, Humphrey TJ. Role of the alternative sigma factors sigmaE and sigmaS in survival of *Salmonella enterica* serovar *Typhimurium* during starvation, refrigeration and osmotic shock. *Microbiology* 2007; **153(Pt 1)**: 263-9.

Payne JB, Osborne JA, Jenkins PK, Sheldon BW. Modeling the growth and death kinetics of *Salmonella* in poultry litter as a function of pH and water activity. *Poult Sci* 2007; **86(1)**: 191-201.

Nde CW, McEvoy JM, Sherwood JS, Logue CM. Cross contamination of turkey carcasses by *Salmonella* species during defeathering. *Poult Sci* 2007; **86(1)**: 162-7.

Lues JF, Theron MM, Venter P, Rasephei MH. Microbial composition in bioaerosols of a high-throughput chicken-slaughtering facility. *Poult Sci* 2007; **86(1)**: 142-9.

Bak H, Ekeroth L, Houe H. Quality control using a multilevel logistic model for the Danish pig *Salmonella* surveillance antibody-ELISA programme. *Prev Vet Med* 2007; **78(2)**: 130-41.

Tabak M, Scher K, Hartog E, Romling U, Matthews KR, Chikindas ML, Yaron S. Effect of triclosan on *Salmonella typhimurium* at different growth stages and in biofilms. *FEMS Microbiol Lett* 2006 Dec 6. Epub ahead of print.

Coburn B, Grassl GA, Finlay BB. *Salmonella*, the host and disease: a brief review. *Immunol Cell Biol* 2006 Dec 5. Epub ahead of print.

Jones TF, Ingram LA, Fullerton KE, Marcus R, Anderson BJ, McCarthy PV, Vugia D, Shiferaw B, Haubert N, Wedel S, Angulo FJ. A case-control study of the epidemiology of sporadic *Salmonella* infection in infants. *Pediatrics* 2006; **118(6)**: 2380-7.

McMahon MA, Xu J, Moore JE, Blair IS, McDowell DA. Environmental stress and antibiotic resistance in food-related pathogens. *Appl Environ Microbiol* 2007; **73(1)**: 211-7.

Thorrold CA, Letsoalo ME, Duse AG, Marais E. Efflux pump activity in flouoroquinolone and tetracycline resistant *Salmonella* and *E. coli* implicated in reduced susceptibility to household antimicrobial cleaning agents. *Int J Food Microbiol* 2007; **113(3)**: 315-20.

Kostic T, Weilharter A, Rubino S, Delogu G, Uzzau S, Rudi K, Sessitch A, Bodrossy L. A microbial diagnostic microarray technique for the sensitive detection and identification of pathogenic bacteria in a background of nonpathogens. *Annal Biochem* 2007; **360(2)**: 244-54.

de Jong P, Vissers MM, van der Meer R, Bovee-Oudenhoven IM. In silico model as a tool for interpretation of intestinal infection studies. *Appl Environ Microbiol* 2007; **73(2)**: 508-15.

Bauer-Garland J, Frye JG, Gray JT, Berrang ME, Harrison MA, Fedorka-Cray PJ. Transmission of *Salmonella enterica* serotype *Typhimurium* in poultry with and without antimicrobial selective pressure. *J Appl Microbiol* 2006; **101(6)**: 1301-8.

Schotte U, Borchers D, Wulff C, Geue L. *Salmonella Montevideo* outbreak in military kennel dogs caused by contaminated commercial feed, which was only recognized through monitoring. *Vet Microbiol* 2007; **119(2-4)**: 316-23.

Warnick LD, Nielsen LR, Nielsen J, Griener M. Simulation model estimates of test accuracy and predictive values for the Danish *Salmonella* surveillance program in dairy herds. *Prev Vet Med* 2006; **77(3-4)**: 284-303.

- Goldbach SG, Alban L. A cost-benefit analysis of *Salmonella*-control strategies in Danish pork production. *Prev Vet Med* 2006; **77(1-2)**: 1-14.
- De Reu K, Grijspeerdt K, Messens W, Heyndrickx M, Uyttendaele M, Debevere J, Herman L. Eggshell factors influencing eggshell penetration and whole egg contamination by different bacteria, including *Salmonella enteridis*. *Int J Food Microbiol* 2006; **112(3)**: 253-60.
- Marcus R, Varma JK, Medus C, Boothe EJ, Anderson BJ, Crume T, Fullerton KE, Moore MR, White PL, Lyszkowicz E, Voetsch AC, Angulo FJ; Emerging Infections Program FoodNet Working Group. Re-assessment of risk factors for sporadic *Salmonella* serotype *Enteritidis* infections: a case-control study in five FoodNet Sites, 2002-2003. *Epidemiol Infect* 2007; **135(1)**: 84-92.
- Gradel KO, Dethlefsen C, C Schonheyder H, Ejlersen T, T Sorensen H, Thomsen RW, Nielsen H. Severity of infection and seasonal variation of non-typhoid *Salmonella* occurrence in humans. *Epidemiol Infect* 2007; **135(1)**: 93-9.
- Van de Giessen A W, Bouwknecht M, Dam-Deisz WD, van Pelt W, Wannet WJ, Visser G. Surveillance of *Salmonella* spp. and *Campylobacter* spp. in poultry production flocks in The Netherlands. *Epidemiol Infect* 2006; **134(6)**: 266-75.
- Galanakis E, Bitsori M, Maraki S, Giannakopolou C, Samonis G, Tselentis Y. Invasive non-typhoidal salmonellosis in immunocompetent infants and children. *Int J Infect Dis* 2007; **11(1)**: 36-9.
- Hughes C, Gillespie IA, O'Brien SJ. Foodborne transmission of infectious intestinal disease in England and Wales 1992-2003. *Food Control* 2007; **18(7)**: 766-772.
- Juneja VK, Melendres MV, Huang L, Gumudavelli V, Subbiah J, Thippareddi H. Modeling the effect of temperature on growth of *Salmonella* in chicken. *Food Microbiol* 2007; **24(4)**: 328-335.
- Jackson V, Blair IS, McDowell DA, Kennedy J, Bolton DJ. The incidence of significant foodborne pathogens in domestic refrigerators. *Food Control* 2007; **18(4)**: 346-351.
- Ukuku DO, Sapers GM, Effect of time before storage and storage temperature on survival of *Salmonella* inoculated on fresh-cut melons. *Food Microbiol* 2007; **24(3)**: 288-295.
- Ou D, Mittal GS. Single-sided pan frying of frozen hamburgers with flippings for microbial safety using modelling and simulation. *Journal of Food Engineering* 2007; **80(1)**: 33-45.

Staphylococcus aureus

Lues JF, Theron MM, van Tonder I. Food-associated bacteria in bioaerosols of delicatessens. *Int J Environ Health Res* 2006; **16(6)**: 419-26

Benedsgaard TW, Thamsborg SM, Aarestrup FM, Enevoldsen C, Vaarst M, Christoffersen AB. Resistance to penicillin of *Staphylococcus aureus* isolates from cows with high somatic cell counts in organic and conventional dairy herds in Denmark. *Acta Vet Scand* 2006; **48(1)**: 24.

Hassani M, Cebrian G, Manas P, Condon S, Pagan R. Induced thermotolerance under nonisothermal treatments of a heat sensitive and a resistant strain of *Staphylococcus aureus* in media of different pH. *Lett Appl Microbiol* 2006; **43(6)**: 619-24.

Soejema T, Nagao E, Yano Y, Yamagata H, Kagi H, Shinagawa K. Risk evaluation for staphylococcal food poisoning in processed milk produced with skim milk powder. *Int J Food Microbiol* 2006 Nov 22. Epub ahead of print.

Other

Steinmuller N, Demma L, Bender JB, Eidson M, Angulo FJ. Outbreaks of enteric disease associated with animal contact: not just a foodborne problem anymore. *Clin Infect Dis* 2006; **43(12)**: 1596-602.

Jones TF, Angulo FJ. Eating in restaurants: a risk factor for foodborne disease? *Clin Infect Dis* 2006; **43(10)**: 1324-8.