

ADVISORY COMMITTEE ON THE MICROBIOLOGICAL SAFETY OF FOOD

Items of possible interest from the literature

A list of items from the literature which may be of interest to Members is attached.

**Secretariat
November 2004**

Campylobacter

Oyarzabal OA, Hawk C, Bilgili SF, Warf CC, Kemp GK. Effects of postchill application of acidified sodium chlorate to control *Campylobacter* spp. and *Escherichia coli* on commercial broiler carcasses. J Food Prot. 2004; **67(10)** : 2288-91

Vandenberg O. Arcobacter species in humans. Emerg Infect Dis. 2004; **10(10)**:1863-7.

Hald B, Skovgard H, Bang DD, Pedersen K, Dybdahl J, Jespersen JB, Madsen M. Flies and *Campylobacter* infection of broiler flocks. Emerg Infect Dis. 2004; **10(8)** :1490-2.

Taboada EN, Acedillo RR, Carrillo CD, Findlay WA, Medeiros DT, Mykytczuk OL, Roberts MJ, Valencia CA, Farber JM, Nash JH. Large-scale comparative genomics meta-analysis of *Campylobacter jejuni* isolates reveals low level of genome plasticity. J Clin Microbiol. 2004; **42(10)** :4566-76.

Ekdahl K, Giesecke J. Travellers returning to Sweden as sentinels for comparative disease incidence in other European countries, *Campylobacter* and *giardia* infection as examples. Euro Surveill. 2004; **9(9)**.

Nicholson FA, Groves SJ, Chambers BJ. Pathogen survival during livestock manure storage and following land application. Bioresour Technol. 2005; **96(2)** :135-43.

Bywater R, Deluyker H, Deroover E, de Jong A, Marion H, McConville M, Rowan T, Shryock T, Shuster D, Thomas V, Valle M, Walters J. A European survey of antimicrobial susceptibility among zoonotic and commensal bacteria isolated from food-producing animals. J Antimicrob Chemother. 2004; **54(4)**:744-54.

Butzler JP. *Campylobacter*, from obscurity to celebrity. Clin Microbiol Infect. 2004; **10(10)** :868-76.

Hansson I, Engvall EO, Lindblad J, Gunnarsson A, Vagsholm I. Surveillance programme for *Campylobacter* species in Swedish broilers, July 2001 to June 2002. Vet Rec. 2004; **155(7)** :193-6.

Ronner AC, Engvall EO, Andersson L, Kaijser B. Species identification by genotyping and determination of antibiotic resistance in *Campylobacter jejuni* and *Campylobacter coli* from humans and chickens in Sweden. Int J Food Microbiol. 2004; **96(2)** :173-9.

Hutchison ML, Walters LD, Moore A, Crookes KM, Avery SM. Effect of length of time before incorporation survival of pathogenic bacteria present in livestock wastes applied to agricultural soil. Appl Environ Microbiol. 2004; **70(9)** :5111-8.

Miller G, Dunn GM, Smith Palmer A, Ogden ID, Strachan NJ. Human *campylobacteriosis* in Scotland: seasonality, regional trends and bursts of infection. Epidemiol Infect. 2004; **132(4)** :585-93.

Hook H, Ekegren MB, Ericsson H, Vagsholm I, Danielsson-Tham ML. Genetic and epidemiological relationships among *Campylobacter* isolates from humans. Scand J Infect Dis. 2004; **36(6-7)** :435-42.

Ramabu SS, Boxall NS, Madie P, Fenwick SG. Some potential sources for transmission of *Campylobacter jejuni* to broiler chickens. Lett Appl Microbiol. 2004; **39(3)** : 252-6.

Bashor MP, Curtis PA, Keener KM, Sheldon BW, Kathariou S, Osbourne JA. Effects of carcass washers on *Campylobacter* contamination in large broiler processing plants. Poult Sci. 2004; **83(7)** :1232-9.

Berrang ME, Northcutt JK, Cason JA. Recovery of *Campylobacter* from broiler faeces during extended storage of transport cages. Poult Sci. 2004; **83(7)** :1213-7

Whyte P, McGill K, Cowley D, Madden RH, Moran L, Scates P, Carroll C, O'Leary A, Fanning S, Collins JD, McNamara E, Moore JE, Cormican M. Occurrence of *Campylobacter* in retail foods in Ireland. Int J Food Microbiol. 2004; **95(2)** :111-8.

Smith K, Reimers N, Barnes HJ, Lee BC, Siletzky R, Kathariou S. *Campylobacter* colonization of sibling turkey flocks reared under different management conditions. J Food Prot. 2004; **67(7)** :1463-8.

Josefsen MH, Cook N, D'Agostino M, Hansen F, Wagner M, Demnerova K, Heuvelink AE, Tassios PT, Linkmark H, Kmet V, Barbanera M, Fach P, Loncarevic S, Hoorfar J. Validation of a PCR-based method for detection of food-borne thermotolerant *campylobacters* in a multicenter collaborative trial. Appl Environ Microbiol. 2004; **70(7)** :4379-83.

Connerton PL, Loc Carrillo CM, Swift C, Dillon E, Scott A, Rees CE, Dodd CE, Frost J, Connerton IF. Longitudinal study of *Campylobacter jejuni* bacteriophages and their hosts from broiler chickens. Appl Environ Microbiol. 2004; **70(7)**: 3877-83.

Clostridium perfringens

Smedley JG 3rd, Fisher DJ, Sayeed S, Chakrabarti G, McClane BA. The enteric toxins of *Clostridium perfringens*. Rev Physiol Biochem Pharmacol. 2004 Oct 23 [Epub ahead of print].

McCabe-Sellers BJ, Beattie SE. Food safety: Emerging trends in foodborne illness surveillance and prevention. J Am Diet Assoc. 2004; **104(11)** :1708-17.

Sabah JR, Juneja VK, Fung DY. Effect of spices and organic acids on the growth of *Clostridium perfringens* during cooling of cooked ground beef. J Food Prot. 2004; **67(9)** :1840-7.

Garrido MN, Skjervheim M, Oppegaard H, Sorum H. Acidified litter benefits the intestinal flora balance of broiler chickens. Appl Environ Microbiol. 2004; **70(9)**:5208-13.

E coli O157

Gupta A. *Escherichia coli* O157 Cluster Evaluation. Emerg Infect Dis. 2004; **10(10)** :1856-8.

Robinson SE, Wright EJ, Hart CA, Bennett M, French NP. Intermittent and persistent shedding of *Escherichia coli* O157 in cohorts of naturally infected calves. J Appl Microbiol. 2004; **97(5)** :1045-53.

Zhao X, Hilliard LR, Mechery SJ, Wang Y, Bagwe RP, Jin S, Tan W. A rapid bioassay for single bacterial cell quantitation using bioconjugated nanoparticles. Proc Natl Acad Sci U S A. 2004; **101(42)** :15027-32.

Silvestro L, Caputo M, Blancato S, Decastelli L, Fioravanti A, Tozzoli R, Morabito S, Caprioli A. Asymptomatic carriage of verocytotoxin-producing *Escherichia coli* O157 in farm workers in Northern Italy. Epidemiol Infect. 2004; **132(5)** :915-9.

Pearce MC, Fenlon D, Low JC, Smith AW, Knight HI, Evans J, Foster G, Synge BA, Gunn GJ. Distribution of *Escherichia coli* O157 in bovine fecal pats and its impact on estimates of the prevalence of fecal shedding. Appl Environ Microbiol. 2004; **70(10)** :5737-43.

Moxley RA. *Escherichia coli* O157:H7: an update on intestinal colonization and virulence mechanisms. Amin Health Res Rev. 2004; **5(1)** :15-33.

Bonardi S, Foni E, Brindani F, Bacci C, Chiapponi C, Cavallini P. Detection and characterization of verocytotoxin-producing *Escherichia coli* (vtec) O157 and non-O157 in cattle at slaughter. New Microbiol. 2004; **27(3)** :255-61.

Ebel E, Schlosser W, Kause J, Orloski K, Roberts T, Narrod C, Malcolm S, Coleman M, Powell M. Draft risk assessment of the public health impact of *Escherichia coli* O157:H7 in ground beef. J Food Prot. 2004; **67(9)** :1991-9.

Baylis CL, MacPhee S, Robinson AJ, Griffiths R, Lilley K, Betts RP. Survival of *Escherichia coli* O157:H7, O111:H- and O26:H11 in artificially contaminated chocolate and confectionery products. Int J Food Microbiol. 2004; **96(1)** :35-48.

Cornick NA, Helgerson AF. Transmission and infectious dose of *Escherichia coli* O157:H7 in swine. Appl Environ Microbiol. 2004; **70(9)** :5331-5.

Huang L. Thermal resistance of *Listeria monocytogenes*, *Salmonella Heidelberg*, and *Escherichia coli* O157:H7 at elevated temperatures. J Food Prot. 2004; **67(8)** :1666-70.

Listeria monocytogenes

Somers EB, Wong AC. Efficacy of two cleaning and sanitizing combinations on *Listeria monocytogenes* biofilms formed at low temperature on a variety of

materials in the presence of ready-to-eat meat residue. J Food Prot. 2004; **67(10)** :2218-29.

Legan JD, Seman DL, Milkowski AL, Hirschey JA, Vandeven MH. Modelling the growth boundary of *Listeria monocytogenes* in ready-to-eat cooked meat products as a function of the product salt, moisture, potassium lactate, and sodium diacetate concentrations. J Food Prot. 2004; **67(10)** :2195-204.

Caggia C, Randazzo CL, Di Salvo M, Romeo F, Giudici P. Occurrence of *Listeria monocytogenes* in green table olives. J Food Prot. 2004; **67(10)** :2189-94.

Dussurget O, Pizarro-Cerda J, Cossart P. Molecular determinants of *Listeria monocytogenes* virulence. Annu Rev Microbiol. 2004; **58** :587-610.

Shin JH, Chang S, Kang DH. Application of antimicrobial ice for reduction of foodborne pathogens (*Escherichia coli* O157:H7, *Salmonella Typhimurium*, *Listeria monocytogenes*) on the surface of fish. J Appl Microbiol. 2004; **97(5)**:916-22.

Salvat G, Fravola P. Risk assessment strategies for Europe: integrated safety strategy or final product control: Example of *Listeria monocytogenes* in processed products from pork meat industry. Dtsch Tierarztl Wochenschr. 2004; **111(8)** :331-4.

Gray MJ, Zadoks RN, Fortes ED, Dogan B, Cai S, Chen Y, Scott VN, Gombas DE, Boor KJ, Wiedmann M. *Listeria monocytogenes* isolates from foods and humans form distinct but overlapping populations. Appl Environ Microbiol. 2004; **70(10)** :5833-41.

Kabuki DY, Kuaye AY, Wiedmann M, Boor KJ. Molecular Subtyping and Tracking of *Listeria monocytogenes* in Latin-Style Fresh-Cheese Processing Plants. J Dairy Sci. 2004; **87(9)** :2803-12.

Cabrita P, Correia S, Ferreira-Dias S, Brito L. Genetic characterization of *Listeria monocytogenes* food isolates and pathogenic potential within serovars 1/2a and 1/2b. Syst Appl Microbiol. 2004; **27(4)** :454-61.

Stecchini ML, Del Torre M, Venir E. Growth of *Listeria monocytogenes* as influenced by viscosity and water activity. Int J Food Microbiol. 2004; **96(2)**:181-7.

Heir E, Lindstedt BA, Rotterud OJ, Vardund T, Kapperud G, Nesbakken T. Molecular epidemiology and disinfectant susceptibility of *Listeria monocytogenes* from meat processing plants and human infections. Int J Food Microbiol. 2004; **96(1)** :85-96.

Rosset P, Cornu M, Noel V, Morelli E, Poumeyrol G. Time-temperature profiles of chilled ready-to-eat foods in school catering and probabilistic analysis of *Listeria monocytogenes* growth. Int J Food Microbiol. 2004; **96(1)** :49-59.

Hayman MM, Baxter I, O'Riordan PJ, Stewart CM. Effects of high-pressure processing on the safety, quality and shelf life of ready-to-eat meats. J Food Prot. 2004; **67(8)** :1709-18.

Jiang X, Islam M, Morgan J, Doyle MP. Fate of *Listeria monocytogenes* in bovine manure-amended soil. J Food Prot. 2004; **67(8)** :1676-81.

Norovirus

De Medici D, Croci L, Suffredini E, Toti L. Reverse transcription-booster PCR for detection of noroviruses in shellfish. Appl Environ Microbiol. 2004; **70(10)**:6329-32.

Radford AD, Gaskell RM, Hart CA. Human norovirus infection and the lessons from animal caliciviruses. Curr Opin Infect Dis. 2004; **17(5)** :471-8.

Barker J, Vipond IB, Bloomfield SF. Effects of cleaning and disinfection in reducing the spread of Norovirus contamination via environmental surfaces. J Hosp Infect. 2004; **58(1)** :42-9.

Hewitt J, Greening GE. Survival and persistence of norovirus, hepatitis A virus, and feline calicivirus in marinated mussels. J Food Prot. 2004; **67(8)** :1743-50.

Pommepuy M, Dumas F, Caprais MP, Camus P, Le Mennec C, Parnaudeau S, Haugarreau L, Sarrette B, Vilagines P, Pothier P, Kholi E, Le Guyader F. Sewage impact on shellfish microbial contamination. Water Sci Technol. 2004; **50(1)** :117-24.

Ueki Y, Akiyama K, Watanabe T, Omura T. Genetic analysis of noroviruses taken from gastroenteritis patients, river water and oysters. Water Sci Technol. 2004; **50(1)** :51-6.

Salmonella

Kankwatira AM, Mwafulirwa GA, Gordon MA. Non-typhoidal *salmonella* bacteraemia—an under-recognized feature of AIDS in African adults. Trop Doct. 2004; **34(4)** :198-200.

Gradel KO, Sayers AR, Davies RH. Surface disinfection tests with *Salmonella* and a putative indicator bacterium, mimicking worst-case scenarios in poultry houses. Poult Sci. 2004; **83(10)** :1636-43.

Sivapalasingham S, Friedman CR, Cohen L, Tauxe RV. Fresh produce: a growing cause of outbreaks of foodborne illness in the United States, 1973 through 1997. J Food Prot. 2004; **67(10)** :2342-53.

Nummer BA, Harrison JA, Harrison MA, Kendall P, Sofos JN, Andress EL. Effects of preparation methods on the microbiological safety of home-dried meat jerky. J Food Prot. 2004; **67(10)** :2337-41.

Baron F, Briandet R, Lesne J, Humbert F, Ablain W, Gautier M. Influence of a nonfavorable environment, egg white, on resistance to heat and disinfectant, adhesion, and virulence of *Salmonella enteritidis*. J Food Prot. 2004; **67(10)**:2269-73.

Mazurek J, Salehi E, Propes D, Holt J, Bannerman T, Nicholson LM, Bundesen M, Duffy R, Moolenaar RL. A multistate outbreak of *Salmonella enterica* serotype *typhimurium* infection linked to raw milk consumption—Ohio, 2003. *J Food Prot.* 2004; **67(10)** :2165-70.

Berghold C, Kornschöber C, Lederer I, Allerberger F. Occurrence of *Salmonella Enteritidis* phage type 29 in Austria: an opportunity to assess the relevance of chicken meat as source of human *salmonella* infections. *Euro Surveill.* 2004; **9(10)** [Epub ahead of print].

Mumma GA. Egg Quality Assurance Programs and Egg-associated *Salmonella Enteritidis* Infections, United States. *Emerg Infect Dis.* 2004; **10(10)** :1782-9.

McCabe-Sellers BJ, Beattie SE. Food safety: Emerging trends in foodborne illness surveillance and prevention. *J Am Diet Assoc.* 2004; **104(11)** :1708-17.

Hensel M. Evolution of pathogenicity islands of *Salmonella enterica*. *Int J Med Microbiol.* 2004; **294(2-3)** :95-102.

O'Hare C, Doran G, Delappe N, Morris D, Buckley V, Corbett-Feeney G, McKeown P, Anderson W, Cormican M. Antimicrobial resistance and phage types of human and non-human *Salmonella enterica* isolates in Ireland, 1998-2003. *Commun Dis Public Health.* 2004; **7(3)** :193-9.

Little CL, Mitchell RT; Food Standards Agency; Local Authorities Coordinators of Regulatory Services; Health Protection Agency. Microbiological quality of pre-cut fruit, sprouted seeds, and unpasteurised fruit and vegetable juices from retail and production premises in the UK, and the application of HACCP. *Commun Dis Public Health.* 2004; **7(3)** :184-90.

Helms M, Simonsen J, Molbak K. Quinolone resistance is associated with increased risk of invasive illness or death during infection with *Salmonella* serotype *Typhimurium*. *J Infect Dis.* 2004; **190(9)** :1652-4.

Guerin PJ, De Jong B, Heir E, Hasseltvedt V, Kapperud G, Styrmö K, Gondrosen B, Lassen J, Andersson Y, Aavitsland P. Outbreak of *Salmonella Livingstone* infection in Norway and Sweden due to contaminated processed fish products. *Epidemiol Infect.* 2004; **132(5)** :889-95.

Bremer V, Leitmeyer K, Jensen E, Metzger U, Meczulat H, Weise E, Werber D, Tschaeppe H, Kreienbrock L, Glaser S, Ammon A. Outbreak of *Salmonella Goldcoast* infections linked to consumption of fermented sausage, Germany 2001. *Epidemiol Infect.* 2004; **132(5)** :881-7.

Salvat G, Fravallo P. Risk assessment strategies for Europe: integrated safety strategy or final product control: Example of *Listeria monocytogenes* in processed products from pork meat industry. *Dtsch Tierärztl Wochenschr.* 2004; **111(8)** :331-4.

Dalton CB, Gregory J, Kirk MD, Stafford RJ, Givney R, Kraa E, Gould D. Foodborne disease outbreaks in Australia, 1995 to 2000. *Commun Dis Intell.* 2004; **28(2)** :211-24. Review.

Evers EG. Predicted quantitative effect of logistic slaughter on microbial prevalence. *Prev Vet Med.* 2004; **65(1-2)** :31-46.

Gardner IA. An epidemiologic critique of current microbial risk assessment practices: the importance of prevalence and test accuracy data. *J Food Prot.* 2004; **67(9)** :2000-7.

Kusumaningrum HD, van Asselt ED, Beumer RR, Zwietering MH. A quantitative analysis of cross-contamination of *Salmonella* and *Campylobacter* spp. via domestic kitchen surfaces. *J Food Prot.* 2004; **67(9)** :1892-903.