

ADVISORY COMMITTEE ON THE MICROBIOLOGICAL SAFETY OF FOOD

Items of possible interest from the literature

A list of items from the literature which may be of interest to members is attached.

**Secretariat
June 2006**

Bacillus cereus

Rajkovic A, Uyttendaele M, Ombregt SA, Jaaskelainen E, Salkinoja-Salonen M, Debevere J. Influence of type of food on the kinetics and overall production of *Bacillus cereus* emetic toxin. J Food Prot 2006; **69(4)**: 847-52.

Han CS, Xie G, Challacombe JF, Altherr MR, Bhotika SS, Bruce D, Campbell CS, Campbell ML, Chen J, Chertkov O, Cleland C, Dimitrijevic M, Doggett NA, Fawcett JJ, Glavina T, Goodwin LA, Hill KK, Hitchcock P, Jackson PJ, Keim P, Kewalramani AR, Longmire J, Lucas S, Malfatti S, McMurry K, Meincke LJ, Misra M, Moseman BL, Mundt M, Munk AC, Okinaka RT, Parson-Quintana B, Reilly LP, Richardson P, Robinson DL, Rubin E, Saunders E, Tapia R, Tesmer JG, Thayer N, Thompson LS, Tice H, Ticknor LO, Wills PL, Brettin TS, Gilna P. Pathogenic sequence analysis of *Bacillus cereus* and *Bacillus thuringiensis* isolates closely related to *Bacillus anthracis*. J Bacteriol 2006; **188(9)**: 3382-90

Moravek M, Dietrich R, Buerk C, Broussolle V, Guidebretiere MH, Granum PE, Nguyen-The C, Martlbauer E. Determination of the toxic potential of *Bacillus cereus* isolates by quantitative enterotoxin analyses. FEMS Microbiol Lett 2006; **257(2)**: 293-298.

Klee SR, Nattermann H, Becker S, Urban-Schriefer M, Franz T, Jacob D, Appel B. Evaluation of different methods to discriminate *Bacillus anthracis* from other bacteria of the *Bacillus cereus* group. J Appl Microbiol 2006; **100(4)**: 673-81.

Carlin F, Fricker M, Pielaat A, Heisterkamp S, Shaheen R, Salkinoja Salonen M, Svensson B, Nguyen-The C, Ehling-Schulz M. Emetic toxin-producing strains of *Bacillus cereus* show distinct characteristics within the *Bacillus cereus* group. Int J Food Microbiol 2006 Feb 22 [Epub ahead of print]

Sorokin A, Candelon B, Guilloux K, Galleron N, Wackerow-Kouzova N, Ehrlich SD, Bourguet D, Sanchis V. Multiple-locus sequence typing analysis of *Bacillus cereus* and *Bacillus thuringiensis* reveals separate clustering and a distinct population structure of psychrotrophic strains. Appl Environ Microbiol 2006; **72(2)**: 1569-78.

Shaheen R, Andersson MA, Apetroaie C, Schulz A, Ehling-Schulz M, Ollilainen VM, Salkinoja-Salonen MS. Potential of selected infant food formulas for production of *Bacillus cereus* emetic toxin, cereulide. Int J Microbiol 2006; **107(3)**: 287-94.

Membre JM, Amezcua A, Bassett J, Giavedoni P, Blackburn Cde W, Gorris LG. A probabilistic modeling approach in thermal inactivation: estimation of postprocess *Bacillus cereus* spore prevalence and concentration. J Food Prot 2006; **69(1)**: 118-29.

Altayar M, Sutherland AD. *Bacillus cereus* is common in the environment but emetic toxin producing isolates are rare. J Appl Microbiol 2006; **100(1)**: 7-14.

Collado J, Fernandex A, Rodrigo M, Martinez A. Modelling the effect of a heat shock and germinant concentration on spore germination of a wild strain of *Bacillus cereus*. Int J Food Microbiol 2006; **106(1)**: 85-9.

Gaillard S, Leguerinel I, Savy N, Mafart P. Quantifying the combined effects of the heating time, the temperature and the recovery medium pH on the regrowth lag time of *Bacillus cereus* spores after a heat treatment. Int J Food Microbiol 2005; **105(1)**: 53-8.

Campylobacter

Idris U, Lu J, Maier M, Sanchez S, Hofacre CL, Harmon BG, Maurer JJ, Lee MD. Dissemination of Fluoroquinolone-Resistant *Campylobacter* spp. within an Integrated Commercial Poultry Production System. Appl Environ Microbiol 2006; **72(5)**: 3441-7.

Zheng J, Meg J, Zhao S, Singh R, Song W. Adherence to and invasion of human intestinal epithelial cells by *Campylobacter jejuni* and *Campylobacter coli* isolates from retail meat products. J Food Prot 2006; **69(4)**: 768-74.

Scherer K, Bartelt E, Sommerfeld C, Hildebrandt G, Quantification of *Campylobacter* on the surface and in the muscle of chicken legs at retail. J Food Prot 2006; **69(4)**: 757-61.

Funakoshi K, Koga M, Takahashi M, Hirata K, Yuki N. *Campylobacter coli* enteritis and Guillain-Barre syndrome: No evidence of molecular mimicry and serological relationship. J Neurol Sci 2006 Apr 4; [Epub ahead of print]

Acik MN, Cetinkaya B. Heterogeneity of *Campylobacter jejuni* and *Campylobacter coli* strains from healthy sheep. Vet Microbiol 2006 Mar 27; [Epub ahead of print]

Murphy C, Carroll C, Jordan KN. Environmental survival mechanisms of the foodborne pathogen *Campylobacter jejuni*. J Appl Microbiol 2006; **100(4)**: 623-32

Manfreda G, De Cesare A, Bondioli V, Stern NJ, Franchini A. Enumeration and identity of *Campylobacter* spp. in Italian broilers. Poult Sci 2006; **85(3)**: 556-62.

Girard MP, Steele D, Chaignat Kieny MP. A review of vaccine research and development: human enteric infections. Vaccine 2006; **24(15)**: 2732-50.

Rasschaert G, Houf K, Van Hende J, De Zutter L. *Campylobacter* contamination during poultry slaughter in Belgium. J Food Prot 2006; **69(1)**: 27-33.

Larkin C, Van Donkersgoed C, Mahdi A, Johnson P, McNab B, Odumeru J. Antibiotic resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from hog, beef, and chicken carcass samples from provincially inspected abattoirs in Ontario. J Food Prot 2006; **69(1)**: 22-6.

Ishihara K, Yamamoto T, Satake S, Takayama S, Kubota S, Negishi H, Kojima A, Asait T, Sawada T, Takahashi T, Tamura Y. Comparison of *Campylobacter* isolated from humans and food-producing animals in Japan. J Appl Microbiol 2006; **100(1)**: 153-60.

Nguyen HT, Corry JE, Miles CA. Heat resistance and mechanism of heat inactivation in thermophilic campylobacters. Appl Environ Microbiol 2006; **72(1)**: 908-13.

Miller WG, Englen MD, Kathariou S, Wesley IV, Wang G, Pittenger-Alley L, Siletz RM, Muraoka W, Fedorka-Cray PJ, Mandrell RE. Identification of host-associated alleles by multilocus sequence typing of *Campylobacter coli* strains from food animals. Microbiology 2006; **152(Pt 1)**: 245-55.

Bohaychuk VM, Gensler GE, King RK, Wu JT, McMullen LM. Evaluation of detection methods for screening meat and poultry products for the presence of foodborne pathogens. J Food Prot 2005; **68(12)**: 2637-47.

Grant A J, Coward C, Jones MA, Woodall CA, Barrow PA, Maskell DJ. Signature-tagged transposon mutagenesis studies demonstrate the dynamic nature of cecal colonization of 2-week-old chickens by *Campylobacter jejuni*. Appl Environ Microbiol 2005; **71(12)**: 8031-41.

Bowen A, Fry A, Richards G, Beauchat L. Infections associated with cantaloupe consumption: a public health concern. Epidemiol Infect 2005 Dec 1; 1-11 [Epub ahead of print]

Clostridium

Margosch D, Ehrmann MA, Buckow R, Heinz V, Vogel RF, Ganzle MG. High-Pressure-Mediated Survival of *Clostridium botulinum* and *Bacillus amyloliquefaciens* Endospores at High Temperature. Appl Environ Microbiol 2006; **72(5)**: 3476-81.

Fallani M, Rogottier-Gois L, Aguilera M, Bridonneau C, Collignon A, Edwards CA, Corthier G, Dore J. *Clostridium difficile* and *Clostridium perfringens* species detected in infant faecal microbiota using 16S rRNA targeted probes. J Microbiol Methods 2006; April 26 [Epub ahead of print]

Rivera VR, Gamez FJ, Keener WK, White JA, Poli MA. Rapid detection of *Clostridium botulinum* toxins A, B, E and F in clinical samples, selected food matrices, and buffer using paramagnetic bead-based electrochemiluminescence detection. Anal Biochem 2006 Mar 20 [Epub ahead of print]

Wohlsen T, Bayliss J, Gray B, Bates J, Katouli M. Evaluation of an alternative method for the enumeration and confirmation of *Clostridium perfringens* from treated and untreated sewages. *Lett Appl Microbiol* 2006; **42(5)**: 438-44.

Johansen CH, Bjerrum L, Finster K, Pedersen K. Effects of a *Campylobacter jejuni* infection on the development of the intestinal microflora of broiler chickens. *Poult Sci* 2006; **85(4)**: 579-87.

Delgado S, Suarez A, Mayo B. Identification of dominant bacteria in feces and colonic mucosa from healthy Spanish adults by culturing and by 16S rDNA sequence analysis. *Dig Dis Sci* 2006; **51(4)**: 744-51.

Delgado S, Ruas-Madiedo P, Suarez A, Mayo B. Interindividual differences in microbial counts and biochemical-associated variables in the feces of healthy Spanish adults. *Dig Dis Sci* 2006; **51(4)**: 737-43.

Lindstrom M, Korkeala H. Laboratory diagnostics of botulism. *Clin Microbiol Rev* 2006; **19(2)**: 298-314.

Cawley GC, Talbot NL, Janacek GJ, Peck MW. Sparse Bayesian kernel survival analysis for modeling the growth domain of microbial pathogens. *IEEE Trans Neural Netw* 2006; **17(2)**: 471-81.

Zanon P, Pattis P, Pittscheider W, Roscia G, De Giorgi G, Sacco G, Botter K, Stockner I, De Giorgi F, Wiedermann CJ. Two cases of foodborne botulism with home-preserved asparagus. *Anaesthesiol Intensivmed Notfallmed Schmerzther* 2006; **41(3)**: 156-9.

Novak J, Katz JA. Probiotics and prebiotics for gastrointestinal infections. *Curr Infect Dis Rep* 2006; **8(2)**: 103-9.

Lindstrom M, Kiviniemi K, Korkeala H. Hazard and control of group II (non-proteolytic) *Clostridium botulinum* in modern food processing. *Int J Food Microbiol* 2006; **108(1)**: 92-104.

Sharma SK, Ferreira JL, Eblen BS, Whiting RC. Detection of type A, B, E, and F *Clostridium botulinum* neurotoxins in foods by using an amplified enzyme-linked immunosorbent assay with digoxigenin-labeled antibodies. *Appl Environ Microbiol* 2006; **72(2)**: 1231-8.

Mueller S, Saunier K, Hanisch C, Norin E, Alm L, Midtvedt T, Cresci A, Silvi S, Orpianesi C, Verenelli MC, Clavel T, Koebnick C, Zunft HJ, Dore J, Blaut M. Differences in fecal microbiota in different European study populations in relation to age, gender, and country: a cross-sectional study. *Appl Environ Microbiol* 2006; **72(2)**: 1027-33.

Chaffer M, Baum M, Grinberg K, Molad T, Elad D. Application of PCR for detection of *Clostridium botulinum* type D in bovine samples. *J Vet Med B Infect Dis Vet Public Health* 2006; **53(1)**: 45-7.

Membre JM, Amezcua A, Bassett J, Giavedoni P, Blackburn Cde W, Gorris LG. A probabilistic modeling approach in thermal inactivation: estimation of postprocess *Bacillus cereus* spore prevalence and concentration. J Food Prot 2006; **69(1)**: 118-29.

Olds DA, Mendonca AF, Sneed J, Bisha B. Influence of four retail food service cooling methods on the behaviour of *Clostridium perfringens* ATCC 10388 in turkey roasts following heating to an internal temperature of 74 degrees C. J Food Prot 2006; **69(1)**: 112-7.

Zhao L, Montville T J, Schaffner DW. Evidence for quorum sensing in *Clostridium botulinum* 56A. Lett Appl Microbiol 2006; **42(1)**: 54-8.

Vernacchio L, Vezina RM, Mitchell AA, Lesko SM, Plaut AG, Acheson DW. Diarrhea in American infants and young children in the community setting: incidence, clinical presentation and microbiology. Pediatr Infect Dis J 2006; **25(1)**: 2-7.

Paredes CJ, Alsaker KV, Papoutsakis ET. A Comparative genomic view of clostridial sporulation and physiology. Nat Rev Microbiol 2005; **3(12)**: 969-78.

Iurlina MO, Fritz R. Characterization of microorganisms in Argentinean honeys from different sources. Int J Food Microbiol 2005; **105(3)**: 297-304.

Nevas M, Lindstrom M, Hautamaki K, Puoskari S, Korkeala H. Prevalence and diversity of *Clostridium botulinum* types A, B, E and F in honey produced in the Nordic countries. Int J Food Microbiol 2005; **105(2)**: 145-51.

Cryptosporidium

Gomez-Couso H, Mendez-Hermida F, Castro-Hermida JA, Ares-Mazas E. Cooking mussels (*Mytilus galloprovincialis*) by steam does not destroy the infectivity of *Cryptosporidium parvum*. J Food Prot 2006; **69(4)**: 948-50.

Smith HV, Caccio SM, Tait A, McLauchlin J, Thompson RC. Tools for investigating the environmental transmission of *Cryptosporidium* and *Giardia* infections in humans. Trends Parasitol 2006; **22(4)**: 160-7.

Li X, Guyot K, Dei-Cas E, Mallard JP, Ballet JJ, Brasseur P. *Cryptosporidium* oocysts in mussels (*Mytilus edulis*) from Normandy (France). Int J Food Microbiol 2006; **108(3)**: 321-5.

Gomez-Couso H, Mendez-Hermida F, Castro-Hermida JA, Ares-Mazas E. *Cryptosporidium* contamination in harvesting areas of bivalve molluscs. J Food Prot 2006; **69(1)**: 185-90.

***E. coli* O157**

Ihekweazu C, Barlow M, Roberts S, Christensen H, Guttridge B, Lewis D, Painter S. Outbreak of *E. coli* O157 infection in the south west of the UK: risks from streams crossing seaside beaches. *Euro Surveill* 2006; **11(4)** [Epub ahead of print].

Walsh C, Duffy G, O'mahony R, Fanning S, Blair IS, McDowell DA. Antimicrobial resistance in Irish isoaltes of verocytotoxigenic *Escherichia coli* (*E. coli*) – VTEC. *Int J Food Microbiol* 2006 April 17 [Epub ahead of print].

Zhang W, Qi W, Albert TJ, Motiwala AS, Alland D, Hyytia-Trees EK, Ribot EM, Fields PI, Whittam TS, Swaminathan B. Probing genomic diversity and evolution of *Escherichia coli* O157 by single-nucleotide polymorphisms. *Genome Res* 2006 Apr 10 [Epub ahead of print].

Rivas M, Miliwebsky E, Chinen I, Roldan CD, Balbi L, Garcia B, Fiorilli G, Sos-Estani S, Kincaid J, Rangel J, Griffin PM; The Case-Control Study Group. Characterization and epideiologic subtyping of Shiga toxin-producing *Escherichia coli* strains isolated from hemolytic uremic syndrome and diarrhea cases in Argentina. *Foodborne Pathog Dis* 2006; **3(1)**: 88-96.

Nielsen EM, Scheutz F, Torpdahl M. Continuous surveillance of Shiga toxin-producing *Escherichia coli* infections by pused-field gel elctrophoresis shows that most infections are sporadic. *Foodborne Pathog Dis* 2006; **3(1)**: 81-7.

Voravuthikunchai SP, Chaowana C, Perepat P, Iida T, Honda T. Antibodies among healthy population of developing countries against enterohaemorrhagic *Escherichia coli* O157: H7. *J Health Popul Nutr* 2005; **23(4)**: 305-10.

Wetzel AN, LeJeune JT. Clonal dissemination of *Escherichia coli* O157: H7 subtypes among dairy farms in northeast Ohio. *Appl Environ Microbiol* 2006; **72(4)**: 2621-6.

Elena SF, Whittam TS, Winkworth CL, Riley MA, Lenski RE. Genomic divergence of *Escherichia coli* strains: evidence for horizontal transfer and variation in mutation rates. *Int Microbiol* 2005; **8(4)**: 271-8.

Shima K, Yoshii N, Akiba M, Nishimura K, Nakazawa M, Yamasaki S. Comparison of PCR-RFLP and PFGE for determining the clonality of enterohemorrhagic *Escherichia coli* strains. *FEMS Microbiol Lett* 2006; **257(1)**: 124-31.

Jordan KN, Maher MM. Sensitive detection of *Escherichia coli* O157:H7 by conventional plating techniques. *J Food Prot* 2006; **69(3)**: 689-92.

Mokhtari A, Frey HC, Jaykus LA. Application of classification and regression trees for sensitivity analysis of the *Escherichia coli* O157: H7 food safety process risk model. J Food Prot 2006; **69(3)**: 609-18.

Moon KD, Delaquis P, Toivonen P, Bach S, Stanich K, Harris L. Destruction of *Escherichia coli* O157:H7 by vanillic acid in unpasteurized juice from six apple cultivars. J Food Prot 2006; **69(3)**: 542-7.

Naugle AL, Holt KG, Levine P, Eckel R. Sustained decrease in the rate of *Escherichia coli* O157: H7-positive raw ground beef samples tested by the food safety and inspection service. J Food Prot 2006; **69(3)**: 480-1.

Raji MA, Minga U, Machangu R. Current epidemiological status of enterohaemorrhagic *Escherichia coli* O157: H7 in Africa. Chin Med J (Engl) 2006; **119(3)**: 217-22.

Jensen C, Ethelberg S, Gervelmeyer A, Nielsen E, Olsen K, Molbak K. First general outbreak of Verocytotoxin-producing *Escherichia coli* O157 in Denmark. Euro Surveill 2006; **11(2)**; [Epub ahead of print].

Foster G, Evans J, Knight HI, Smith AW, Gunn GJ, Allison LJ, Synge BA, Pennycott TW. Analysis of feces samples collected from a wild-bird garden feeding station in Scotland for the presence of verocytotoxin-producing *Escherichia coli* O157. Appl Environ Microbiol 2006; **72(3)**: 2265-7.

Turner J, Bowers RG, Begon M, Robinson SE, French NP. A semi-stochastic model of the transmission of *Escherichia coli* O157 in a typical UK dairy herd: Dynamics, sensitivity analysis and intervention/prevention strategies. J Theor Biol 2006; Mar 1 [Epub ahead of print].

Duffy G, Walsh C, Blair IS, McDowell DA. Survival of antibiotic resistant and antibiotic sensitive strains of *E. coli* O157 and *E. coli* O26 in food matrices. Int J Food Microbiol 2006; Feb 25 [Epub ahead of print].

Lejeune JT, Hancock D, Wasteson Y, Skjerve E, Urdahl AM. Comparison of *E. coli* O157 and Shiga toxin-encoding genes (stx) prevalence between Ohio, USA and Norwegian dairy cattle. Int J Food Microbiol 2006 Feb 24 [Epub ahead of print].

Ejidokun OO, Walsh A, Barnett J, Hope Y, Ellis S, Sharp MW, Paiba GA, Logan M, Willshaw GA, Cheasty T. Human Vero cytotoxigenic *Escherichia coli* (VTEC) O157 infection linked to birds. Epidemiol Infect 2006; **134(2)**: 421-3.

Listeria monocytogenes

Angelidis AS, Koutsoumanis K. Prevalence and concentration of *Listeria monocytogenes* in sliced ready-to-eat meat products in the Hellenic retail market. J Food Prot 2006; **69(4)**: 938-42.

Takeuchi K, Mytle N, Lambert S, Coleman M, Doyle MP, Smith MA. Comparison of *Listeria monocytogenes* virulence in a mouse model. J Food Prot 2006; **69(4)**: 842-6.

Nakamura H, Tokuda Y, Sono A, Koyama T, Ogasawara J, Hase A, Haruki K, Nishikawa Y. Molecular typing to trace *Listeria monocytogenes* isolated from cold-smoked fish to a contamination source in a processing plant. J Food Prot 2006; **69(4)**: 835-41.

Sinigaglia M, Bevilacqua A, Campaniello D, D'Amato D, Corbo MR. Growth of *Listeria monocytogenes* in fresh-cut coconut as affected by storage conditions and inoculum size. J Food Prot 2006; **69(4)**: 820-5.

Gray MJ, Freitag NE, Boor KJ. How the bacterial pathogen *Listeria monocytogenes* mediates the switch from environmental Dr Jekyll to pathogenic Mr Hyde. Infect Immun 2006; **74(5)**: 2505-12.

Czuprynski CJ. *Listeria monocytogenes*: silage, sandwiches and science. Anim Health Res Rev 2005; **6(2)**: 211-7.

Pohl MA, Wiedmann M, Nightingale KK. Associations among *Listeria monocytogenes* genotypes and distinct clinical manifestations of listeriosis in cattle. Am J Vet Res 2006; **67(4)**: 616-26.

Francois K, Kevlieghere F, Uyttendaele M, Standaert AR, Geeraerd AH, Nadal P, Van Impe JF, Debevere J. Single cell variability of *L. monocytogenes* grown on liver pate and cooked ham at 7 degrees C: comparing challenge test data to predictive simulations. J Appl Microbiol 2006; **100(4)**: 800-12.

Vorst KL, Todd EC, Rysert ET. Transfer of *Listeria monocytogenes* during mechanical slicing of turkey breast bologna and salami. J Food Prot 2006; **69(3)**: 619-26.

Jacobsen T, Koch AG. Influence of different histories of the inoculum on lag phase and growth of *Listeria monocytogenes* in meat models. J Food Prot 2006; **69(3)**: 532-41.

Cao J, Clarke M, Witkowsky R, Lu H, Sayedahman A, Levin RE, McLandsborough LA. Concentrations and tracking of *Listeria monocytogenes* strains in a seafood-processing environment using a most-probable-number enrichment procedure and randomly amplified polymorphic DNA analysis. J Food Prot 2006; **69(3)**: 489-94.

[No authors listed] Evaluation of farm management practices as risk factors for clinical listeriosis and fecal shedding of *Listeria monocytogenes* in ruminants. J Am Vet Med Assoc 2006; **228(5)**: 737

Houben JH, Eckenhausen F. Surface pasteurization of vacuum-sealed precooked ready-to-eat meat products. J Food Prot 2006; **69(2)**: 459-68.

Chen Y, Ross WH, Gray MJ, Wiedmann M, Whiting RC, Scott VN. Attributing risk of *Listeria monocytogenes* subgroups: dose response in relation to genetic lineages. J Food Prot 2006; **69(2)**: 335-44.

Francois K, Devlieghere F, Uyttendaele M, Debevere J. Risk assessment of *Listeria monocytogenes*: impact of individual cell variability on the exposure assessment step. Risk Anal 2006; **26(1)**: 105-14.

Yang H, Mokhtari A, Jaykus LA, Morales RA, Cates SC, Cowen P. Consumer phase risk assessment for *Listeria monocytogenes* in deli meats. Risk Anal 2006; **26(1)**: 89-103.

Lin CM, Takeuchi K, Zhang L, Dohm CB, Meyer JD, Hall PA, Doyle MP. Cross-contamination between processing equipment and deli meats by *Listeria monocytogenes*. J Food Prot 2006; **69(1)**: 71-9.

Yoon Y, Skandamis PN, Kendall PA, Smith GC, Sofos JN. A predictive model for the effect of temperature and predrying treatments in reducing *Listeria monocytogenes* populations during drying of beef jerky. J Food Prot 2006; **69(1)**: 62-70.

Millet L, Saubusse M, Didiene R, Tessier L, Montel MC. Control of *Listeria monocytogenes* in raw-milk cheeses. Int J Food Microbiol 2006; **108(1)**: 105-14.

Gilbreth SE, Call JE, Wallace FM, Scott VN, Chen Y, Luchansky JB. Relatedness of *Listeria monocytogenes* isolates recovered from selected ready-to-eat foods and listeriosis patients in the United States. Appl Environ Microbiol 2005; **71(12)**: 8115-22.

Norovirus

Etherington GJ, Ring SM, Charleston MA, Dicks J, Rayward-Smith VJ, Roberts IN. Tracing the origin and co-phylogeny of the caliciviruses. J Gen Virol 2006; **87(Pt 5)**: 1229-35.

Trujillo AA, McCaustland KA, Zheng DP, Hadley LA, Vaughn G, Adams SM, Ando T, Glass RI, Monroe SS. Use of TaqMan real-time reverse transcription-PCR for rapid detection, quantification, and typing of norovirus. J Clin Microbiol 2006; **44(4)**: 1405-12.

Fukuda S, Takao S, Kuwayama M, Shimazu Y, Miyazaki K. Rapid detection of norovirus from fecal specimens by real-time reverse transcription-loop-mediated isothermal amplification assay. J Clin Microbiol 2006; **44(4)**: 1376-81.

Ike AC, Brockmann SO, Hartelt K, Marschang RE, Contzen M, Oehme RM. Molecular epidemiology of norovirus in outbreaks of gastroenteritis in southwest Germany from 2001 to 2004. J Clin Microbiol 2006; **44(4)**: 1262-7.

Waters A, Coughlan S, Dunford L, Hall WW. Molecular epidemiology of Norovirus strains circulating in Ireland from 2003 to 2004. *Epidemiol Infect* 2006; 1-9 [Epub ahead of print].

Malik YS, Goyal SM. Virucidal efficacy of sodium bicarbonate on a food contact surface against feline calicivirus, a norovirus surrogate. *Int J Food Microbiol* 2006; [Epub ahead of print].

Hansman GS, Natori K, Shirato-Horikoshi H, Ogawa S, Oka T, Katayama K, Tanaka T, Miyoshi T, Sakae K, Kobayashi S, Shinohara M, Uchinda K, Sakurai N, Shinozaki K, Okada M, Seto Y, Kamata K, Nagata N, Tanaka K, Miyamura T, Takeda N. Genetic and antigenic diversity among noroviruses. *J Gen Virol* 2006; **87(Pt 4)**: 909-19.

Sasaki Y, Kai A, Hayashi Y, Shinkai T, Noguchi Y, Hasegawa M, Sadamasu K, Mori K, Tabei Y, Nagashima M, Morozumi S, Yamamoto T. Multiple viral infections and genomic divergence among noroviruses during an outbreak of acute gastroenteritis. *J Clin Microbiol* 2006; **44(3)**: 790-7.

Costantini V, Loisy F, Joens L, Le Guyader FS, Saif LJ. Human and animal enteric caliciviruses in oysters from different coastal regions of the United States. *Appl Environ Microbiol* 2006; **72(3)**: 1800-9.

Turcios RM, Widdowson MA, Sulka AC, Mead PS, Glass RI. Reevaluation of epidemiological criteria for identifying outbreaks of acute gastroenteritis due to norovirus: United States, 1998-2000. *Clin Infect Dis* 2006; **42(7)**: 964-9.

Boxman IL, Tilburg JJ, Te Loeke NA, Vennema H, Jonker K, de Boer E, Koopmans M. Detection of noroviruses in shellfish in the Netherlands. *Int J Food Microbiol* 2006; **108(3)**: 391-6.

Wang QH, Han MG, Cheetham S, Souza M, Funk JA, Saif LJ. Porcine noroviruses related to human noroviruses. *Emerg Infect Dis* 2005; **11(12)**: 1874-81.

Malik YS, Maherchandani S, Goyal SM. Comparative efficacy of ethanol and isopropanol against feline calicivirus, a norovirus surrogate. *Am J Infect Control* 2006; **34(1)**: 31-5.

Koopmans M. Food-borne norovirus outbreaks: a nuisance or more than that? *Wien Klin Wochenschr* 2005; **117(23-24)**: 789-91.

Fretz R. [How important is the norovirus relationship to gastroenteritis]. Article in German. *Wien Klin Wochenschr* 2005; **117(23-24)**: 802-8.

Chan MC, Sung JJ, Lam RK, Chan PK, Lai RW, Leung WK. Sapovirus detection by quantitative real-time RT-PCR in clinical stool specimens. *J Virol Methods* 2006; **134(1-2)**: 146-53.

Zheng DP, Ando T, Fankhauser RL, Beard RS, Glass RI, Monroe SS. Norovirus classification and proposed strain nomenclature. *Virology* 2006; **346(2)**: 312-23.

Ng TL, Chan PP, Phua TH, Loh JP, Yip R, Wong C, Liaw CW, Tan BH, Chiew KT, Chua SB, Lim S, Ooi PL, Chew SK, Goh KT. Oyster-associated outbreaks of Norovirus gastroenteritis in Singapore. *J Infect* 2005; **51(5)**: 413-8.

Salmonella

Huang DS, Li DF, Xing JJ, Ma YX, Li ZJ, Lv SQ. Effects of feed particle size and feed form on survival of *Salmonella typhimurium* in the alimentary tract and cecal *S. typhimurium* reduction in growing broilers. *Poult Sci* 2006; **85(5)**: 831-6.

[No authors listed]. Dog and cat foods and treats: a new source of *Salmonella* infections? *Child Health Alert* 2006; **24**: 4.

Van de Giessen AW, Bouwknecht M, Dam-Deisz WD, van Pelt W, Wannet WJ, Visser G. Surveillance of *Salmonella* spp. and *Campylobacter* spp. in poultry production flocks in The Netherlands. *Epidemiol Infect* 2006 May 2: 1-10 [Epub ahead of print].

Kang H, Loui C, Clavijo RI, Riley LW, Lu S. Survival characteristics of *Salmonella enterica* serovar Enteritidis in chicken egg albumen. *Epidemiol Infect* 2006 Feb 28: 1-10 [Epub ahead of print].

O'Connor AM, McKean JD, Beary JH, Brockus SL. Prevalence of exposure to *Salmonella* spp in finishing swine marked in Iowa. *Am J Vet Res* 2006; **67(5)**: 829-33.

Carey-Smith GV, Billington C, Cornelius AJ, Hudson JA, Heinemann JA. Isolation and characterization of bacteriophages infecting *Salmonella* spp. *FEMS Microbiol Lett* 2006; **258(2)**: 182-6.

Forsyth J, Lightfoot D, Powling J, Tan A. *Salmonella* surveillance in Australia – created from adversity. *J AOAC Int* 2006; **89(2)**: 560-5.

Swaminathan B, Barrett TJ, Fields P. Surveillance for human *Salmonella* infections in the United States. *J AOAC Int* 2006; **89(2)**: 553-9.

Lammerding AM. Modeling and risk assessment for *Salmonella* in meat and poultry. *J AOAC Int* 2006; **89(2)**: 543-52.

Cliver DO. Cutting boards in *Salmonella* cross-contamination. *J AOAC Int* 2006; **89(2)**: 538-42.

- Mozola MA. Genetics-based methods for detection of *Salmonella* spp. in foods. J AOAC Int 2006; **89(2)**: 517-29.
- Fletcher DL. Influence of sampling methodology on reported incidence of *Salmonella* in poultry. J AOAC Int 2006; **89(2)**: 512-6.
- Saeed AM, Walk ST, Arshad M, Whittam TS. Clonal structure and variation in virulence of *Salmonella enteritidis* isolated from mice, chickens, and humans. J AOAC Int 2006; **89(2)**: 504-11.
- Poppe C, Martin L, Muckle A, Archambault M, McEwen S, Weir E. Characterization of antimicrobial resistance of *Salmonella* Newport isolated from animals, the environment, and animal food products in Canada. Can J Vet Res 2006; **70(2)**: 105-14.
- Doorduyn Y, Van den Brandhof WE, Van Duynhoven YT, Wannet WJ, Van Pelt W. Risk factors for *Salmonella Enteritidis* and *Typhimurium* (DT104 and non-DT104) infections in The Netherlands: predominant roles for raw eggs in *Enteritidis* and sandboxes in *Typhimurium* infections. Epidemiol Infect 2006; **134(3)**: 617-26.
- Taguchi M, Seto K, Yamazaki W, Tsukamoto T, Izumiya H, Watanabe H. CMY-2 Beta-Lactamase-Producing *Salmonella enterica* serovar *Infantis* isolated from poultry in Japan. Jpn J Infect Dis 2006; **59(2)**: 144-6.
- Meldrum RJ, Smith RM, Wilson IG. Three-year surveillance program examining the prevalence of *Campylobacter* and *Salmonella* in whole retail raw chicken. J Food Prot 2006; **69(4)**: 928-31.
- Nightingale KK, Thippareddi H, Phebus RK, Marsden JL, Nutsch AL. Validation of a traditional Italian-style salami manufacturing process for control of *Salmonella* and *Listeria monocytogenes* J Food Prot 2006; **69(4)**: 794-800.
- Uesugi AR, Harris LJ. Growth of *Salmonella Enteritidis* phage type 30 in almond hull and shell slurries and survival in drying almond hulls. J Food Prot 2006; **69(4)**: 712-8.
- Charlesworth J, Eyre-Walker A. The Rate of Adaptive Evolution in Enteric Bacteria. Mol Biol Evol 2006 Apr 18; [Epub ahead of print].
- Talbot EA, Gagnon ER, Greenblatt J. Common ground for the control of multidrug-resistant *salmonella* in ground beef. Clin Infect Dis 2006; **42(10)**: 1455-62.
- Centers for Disease Control and Prevention (CDC). Preliminary FoodNet data on the incidence of infection with pathogens transmitted commonly through food – 10 States, United States, 2005. MMWR Morb Mortal Wkly Rep 2006; **55(14)**: 392-5.

Lukinmaa S, Nakari UM, Liimatainen A, Siitonen A. Genomic diversity within phage types of *Salmonella enterica* spp. *enterica* serotypes *Enteritidis* and *Typhimurium*. *Foodborne Pathog Dis* 2006; **3(1)**: 97-105.

Fleury M, Charron DF, Holt JD, Allen OB, Maarouf AR. A time series analysis of the relationship of ambient temperature and common bacterial enteric infections in two Canadian provinces. *Int J Biometeorol* 2006 Mar 31; [Epub ahead of print].

Morris D, Whelan M, Corbett-Feeney G, Cormican M, Hawkey P, Li X, Doran G. First report of extended-spectrum-beta-lactamase-producing *Salmonella enterica* isolates in Ireland. *Antimicrob Agents Chemother* 2006; **50(4)**: 1608-9.

McLaughlin JB, Castrodale LJ, Gardner MJ, Ahmed R, Gessner BD. Outbreak of multidrug-resistant *Salmonella typhimurium* associated with ground beef served at a school potluck. *J Food Prot* 2006; **69(3)**: 666-70.

Becker D, Selbach M, Rollenhagen C, Ballmaier M, Meyer TF, Mann M, Bumann D. Robust *Salmonella* metabolism limits possibilities for new antimicrobials. *Nature* 2006; **440(7082)**: 303-7.

Meldrum RJ, Smith RM, Ellis P, Garside J; on behalf of the Welsh Food Microbiological Forum. Microbiological quality of randomly selected ready-to-eat foods sampled between 2003 and 2005 in Wales, UK. *Int J Food Microbiol* 2006; **108(3)**: 397-400.

Srinivasan V, Morowitz HJ. Ancient genes in contemporary persistent microbial pathogens. *Biol Bull* 2006; **210(1)**: 1-9.

Centers for Disease Control and Prevention (CDC). Multistate outbreak of *Salmonella typhimurium* infections associated with eating ground beef – United States, 2004. *MMWR Morb Mortal Wkly Rep* 2006; **55(7)**: 180-2.

Samadpour M, Barbour MW, Nguyen T, Cao TM, Buck F, Depavia GA, Mazengia E, Yang P, Alfi D, Lopes M, Stopforth JD. Incidence of enterohemorrhagic *Escherichia coli*, *Escherichia coli* O157, *Salmonella*, and *Listeria monocytogenes* in retail fresh ground beef, sprouts, and mushrooms. *J Food Prot* 2006; **69(2)**: 441-3.

Pradhan AK, Li Y, Swem BL, Mauromoustakos A. Predictive model for the survival, death, and growth of *Salmonella typhimurium* in broiler hatchery. *Poult Sci* 2005; **84(12)**: 1959-66.

Musgrove MT, Jones DR, Northcutt JK, Harrison MA, Cox NA, Ingram KD, Hinton AJ Jr. Recovery of *Salmonella* from commercial shell eggs by shell rinse and shell crush methodologies. *Poult Sci* 2005; **84(12)**: 1955-8.

Herrera FC, Santos JA, Otero A, Garcia-Lopez ML. Occurrence of foodborne pathogenic bacteria in retail prepackaged portions of marine fish in Spain. *J Appl Microbiol* 2006; **100(3)**: 527-36.

Cobbold RN, Rice DH, Davis MA, Besser TE, Hancock DD. Long-term persistence of multi-drug-resistant *Salmonella enterica* serovar Newport in two dairy herds. *J Am Vet Med Assoc* 2006; **228(4)**: 585-91.

Dechet AM, Scallan E, Gensheimer K, Hoeksstra R, Gunderman-King J, Lockett J, Wrigley D, Chege W, Sobel J; Multistate Working Group. Outbreak of multidrug-resistant *Salmonella enterica* serotype *Typhimurium* Definitive Type 104 infection linked to commercial ground beef, northeastern United States, 2003-2004. *Clin Infect Dis* 2006; **42(6)**: 747-52.

Koehler KM, Lasky T, Fein SB, DeLong SM, Hawkins MA, Rabatsky-Ehr T, Ray SM, Shiferaw B, Swanson E, Vugia DJ. Population-based incidence of infection with selected bacterial enteric pathogens in children younger than five years of age, 1996-1998. *Pediatr Infect Dis J* 2006; **25(2)**: 129-34.

Clavijo RI, Loui C, Andersen GL, Riley LW, Lu S. Identification of genes associated with survival of *Salmonella enterica* serovar *Enteritidis* in chicken egg albumen. *Appl Environ Microbiol* 2006; **72(2)**: 1055-64.

Kosmider R, Kelly L, Evans S, Gettinby G. A statistical system for detecting *Salmonella* outbreaks in British livestock. *Epidemiol Infect* 2006 Jan 26: 1-9 [Epub ahead of print]

Gatto AJ, Peters TM, Green J, Fisher IS, Gill ON, O'Brien SJ, Maguire C, Berghold C, Lederer I, Gerner-Smidt P, Torpdahl M, Siitonen A, Linmaa S, Tschape H, Prager R, Luzzi I, Dionisi AM, Van der Swaluw WK, Heck M, Coia J, Brown D, Usera M, Echeita A, Threlfall EJ. Distribution of molecular subtypes within *Salmonella enterica* serotype *Enteritidis* phage type 3 and *S. Typhimurium* definitive phage type 104 in nine European countries, 2000-2004: results of an international multi-centre study. *Epidemiol Infect* 2006; Jan 25: 1-8 [Epub ahead of print].

Messens W, Grijspeerdt K, Herman L. Eggshell characteristics and penetration by *Salmonella enterica* serovar *Enteritidis* through the production period of a layer flock. *Br Poult Sci* 2005; **46(6)**: 694-700.

Notermans DW, van Pelt W, Kivi M, van de Giessen AW, Wannet WJ, Bosman A. [Sharp increase of infections with *Salmonella* serotype *Typhimurium* DT104 in The Netherlands] [Article in Dutch]. *Ned Tijdschr Geneesk* 2005; **149(53)**: 2992-4.

Vij V, Ailes E, Wolyniak C, Angulo FJ, Klontz KC. Recalls of spices due to bacterial contamination monitored by the US Food and Drug Administration: the predominance of *Salmonellae*. *J Food Prot* 2006; **69(1)**: 233-7.

Papagrigrakis MJ, Yapijakis C, Synodinos PN, Baziotopoulou-Valavani E. DNA examination of ancient dental pulp incriminates typhoid fever as a probably cause of the Plague of Athens. *Int J Infect Dis* 2006; **10(3)**: 206-14.

Oliveira CJ, Carvalho LF, Garcia TB. Experimental airborne transmission of *Salmonella Agona* and *Salmonella Typhimurium* in weaned pigs. *Epidemiol Infect* 2006; **134(1)**: 199-209.

Gebreyes WA, Altier C, Thakur S. Molecular epidemiology and diversity of *Salmonella* serovar *Typhimurium* in pigs using phenotypic and genotypic approaches. *Epidemiol Infect* 2006; **134(1)**: 187-98.

Mackey BM, Kelly AF, Colvin JA, Robbins PT, Fryer PJ. Predicting the thermal inactivation of bacteria in a solid matrix: simulation studies on the relative effects of microbial thermal resistance parameters and process conditions. *Int J Food Microbiol* 2006; **107(3)**: 295-303.

De Jong B, Ekdahl K. The comparative burden of salmonellosis in the European Union members states, associated and candidate countries. *BMC Public Health* 2006; **6**: 4.

Koonse B, Burkhardt W 3rd, Chrtel S, Hoskin GP. *Salmonella* and the sanitary quality of aquacultured shrimp. *J Food Prot* 2005; **68(12)**: 2527-32.

Bowen A, Fry A, Richards G, Beauchat L. Infections associated with cantaloupe consumption: a public health concern. *Epidemiol Infect* 2005 Dec 1: 1-11 [Epub ahead of print].

Davison HC, Smith RP, Pascoe SJ, Sayers AR, Davies RH, Weaver JP, Kidd SA, Dalziel RW, Evans SJ. Prevalence, incidence and geographical distribution of serovars of *Salmonella* on dairy farms in England and Wales. *Vet Rec* 2005; **157(22)**: 703-11.

Stewart WC, Pollock KG, Browning LM, Young D, Smith-Palmer A, Reilly WJ. Survey of zoonoses recorded in Scotland between 1993 and 2002. *Vet Rec* 2005; **157(22)**: 697-702.

Hajmeer M, Basheer I, Hew C, Cliver DO. Modeling the survival of *Salmonella* spp. in chorizos. *Int J Food Microbiol* 2006; **107(1)**: 59-67.

Haimovich B, Venkatesan MM. *Shigella* and *Salmonella*: death as a means of survival. *Microbes Infect* 2006; **8(2)**: 568-77.

Unicomb LE, Simmons G, Merritt T, Gregory J, Nicol C, Jelfs P, Kirk M, Tan A, Thomson R, Adamopoulos J, Little CL, Currie A, Dalton CB. Sesame seed products contaminated with *Salmonella*: three outbreaks associated with tahini. *Epidemiol Infect* 2005; **133(6)**: 1065-72.

Palmgren H, Aspan A, Broman T, Bengtsson K, Blomquist L, Bergstrom S, Sellin M, Wollin R, Olsen B. *Salmonella* in Black-headed gulls (*Larus*

ridibundus); prevalence, genotypes and influence on *Salmonella* epidemiology. *Epidemiol Infect* 2006; **134(3)**: 635-44.

Sumner J, Ross T, Jenson I, Pointon A. A risk microbiological profile of the Australian red meat industry: risk ratings of hazard-product pairings. *Int J Food Microbiol* 2005; **105(2)**: 221-32.

Staphylococcus aureus

Smyth DS, Kennedy J, Twohig J, Miajlovic H, Bolton D, Smyth CJ. *Staphylococcus aureus* isolates from Irish domestic refrigerators possess novel enterotoxin and enterotoxin-like genes and are clonal in nature. *J Food Prot* 2006; **69(3)**: 508-15.

Alarcon B, Vicedo B, Aznar R. PCR-based procedures for detection and quantification of *Staphylococcus aureus* and their application in food. *J Appl Microbiol* 2006; **100(2)**: 352-64.

Berrada H, Soriano JM, Manes J, Pico Y. Real-time quantitative PCR of *Staphylococcus aureus* and application in restaurant meals. *J Food Prot* 2006; **69(1)**: 106-11.

Lalla F, Dingle P, Cheong C. The antibacterial action of cloths and sanitizers and the use of environmental alternatives in food industries. *J Environ Health* 2005; **68(5)**: 31-5.

Bania J, Dabrowska A, Bystron J, Korzekwa K, Chrzanowska J, Molenda J. Distribution of newly described enterotoxin-like genes in *Staphylococcus aureus* from food. *Int J Food Microbiol* 2006; **108(1)**: 36-41.

Jorgensen HJ, Mathisen T, Lovseth A, Omoe K, Qvale KS, Loncarevic S. An outbreak of staphylococcal food poisoning caused by enterotoxin H in mashed potato made with raw milk. *FEMS Microbiol Lett* 2005; **252(2)**: 267-72.