

ADVISORY COMMITTEE ON THE MICROBIOLOGICAL SAFETY OF FOOD
INFORMATION PAPER

Items of possible interest from the literature

Bacillus cereus

Ceuppens S, Uyttendaele M, Drieskens K, Rajkovic A, Boon N, Wiele TV. Survival of *Bacillus cereus* Vegetative Cells and Spores during In Vitro Simulation of Gastric Passage. J Food Prot. 2012 Apr;75(4):690-4.

Guinebretiere MH, Auger S, Galleron N, Contzen M, De Sarrau B, De Buyser ML, Lamberet G, Fagerlund A, Granum PE, Lereclus D, De Vos P, Nguyen-The C, Sorokin A. *Bacillus cytotoxicus* sp. nov. is a new thermotolerant species of the *Bacillus cereus* group occasionally associated with food poisoning. Int J Syst Evol Microbiol. 2012 Feb 17. [Epub ahead of print]

Logan NA. Bacillus and relatives in foodborne illness. J Appl Microbiol. 2012 Mar;112(3):417-29.

Sudhaus N, Pina-Pérez MC, Martínez A, Klein G. Inactivation Kinetics of Spores of *Bacillus cereus* Strains Treated by a Peracetic Acid-Based Disinfectant at Different Concentrations and Temperatures. Foodborne Pathog Dis. 2012 Apr 16. [Epub ahead of print].

Campylobacter

Ahmed MU, Dunn L, Ivanova EP. Evaluation of Current Molecular Approaches for Genotyping of *Campylobacter jejuni* Strains. Foodborne Pathog Dis. 2012 Apr 16. [Epub ahead of print]

Bessell PR, Rotariu O, Innocent GT, Smith-Palmer A, Strachan NJ, Forbes KJ, Cowden JM, Reid SW, Matthews L. Using sequence data to identify alternative routes and risk of infection: A case-study of *Campylobacter* in Scotland. BMC Infect Dis. 2012 Apr 1;12(1):80.

Bullman S, O'Leary J, Corcoran D, Sleator RD, Lucey B. Molecular-based detection of non-culturable and emerging campylobacteria in patients presenting with gastroenteritis. Epidemiol Infect. 2012 Apr;140(4):684-8.

Ellis-Iversen J, Ridley A, Morris V, Sowa A, Harris J, Atterbury R, Sparks N, Allen V. Persistent environmental reservoirs on farms as risk factors for *Campylobacter* in commercial poultry. Epidemiol Infect. 2012 May;140(5):916-24.

Friesema IH, Havelaar AH, Westra PP, Wagenaar JA, van Pelt W. Poultry culling and *Campylobacteriosis* reduction among humans, the Netherlands. *Emerg Infect Dis.* 2012 Mar;18(3):466-8.

Herbinger KH, Drerup L, Alberer M, Nothdurft HD, Sonnenburg Fv, Löscher T. Spectrum of imported infectious diseases among children and adolescents returning from the tropics and subtropics. *J Travel Med.* 2012 May;19(3):150-7.

Liu X, Gao B, Novik V, Galán JE. Quantitative Proteomics of Intracellular *Campylobacter jejuni* Reveals Metabolic Reprogramming. *PLoS Pathog.* 2012 Mar;8(3):e1002562.

Liu Y, Betti M, Gänzle MG. High pressure inactivation of *Escherichia coli*, *Campylobacter jejuni*, and spoilage microbiota on poultry meat. *J Food Prot.* 2012 Mar;75(3):497-503.

Mattheus W, Botteldoorn N, Heylen K, Pochet B, Dierick K. Trend Analysis of Antimicrobial Resistance in *Campylobacter jejuni* and *Campylobacter coli* Isolated from Belgian Pork and Poultry Meat Products Using Surveillance Data of 2004-2009. *Foodborne Pathog Dis.* 2012 Apr 17. [Epub ahead of print]

McCarthy ND, Gillespie IA, Lawson AJ, Richardson J, Neal KR, Hawtin PR, Maiden MC, O'Brien SJ. Molecular epidemiology of human *Campylobacter jejuni* shows association between seasonal and international patterns of disease. *Epidemiol Infect.* 2012 Feb 28:1-9. [Epub ahead of print]

Nielsen HL, Ejlersten T, Engberg J, Nielsen H. High incidence of *Campylobacter concisus* in gastroenteritis in North Jutland, Denmark: a population-based study. *Clin Microbiol Infect.* 2012 Mar 27. doi:10.1111/j.1469-0691.2012.03852.x.

Pittenger LG, Frye JG, McEnerney V, Reeves J, Haro J, Fedorka-Cray PJ, Harrison MA, Englen MD. Analysis of *Campylobacter jejuni* Whole-Genome DNA Microarrays: Significance of Prophage and Hypervariable Regions for Discriminating Isolates. *Foodborne Pathog Dis.* 2012 Apr 16. [Epub ahead of print]

Poli VF, Thorsen L, Olesen I, Wik MT, Jespersen L. Differentiation of the virulence potential of *Campylobacter jejuni* strains by use of gene transcription analysis and a Caco-2 assay. *Int J Food Microbiol.* 2012 Apr 2;155(1-2):60-8.

Powell LF, Lawes JR, Clifton-Hadley FA, Rodgers J, Harris K, Evans SJ, Vidal A. The prevalence of *Campylobacter* spp. in broiler flocks and on broiler carcasses, and the risks associated with highly contaminated carcasses. *Epidemiol Infect.* 2012 Feb 16:1-14.

Sahin O, Fitzgerald C, Stroika S, Zhao S, Sippy RJ, Kwan P, Plummer PJ, Han J, Yaeger MJ, Zhang Q. Molecular evidence for zoonotic transmission of an emergent, highly pathogenic *Campylobacter jejuni* clone in the United States. *J Clin Microbiol.* 2012 Mar;50(3):680-7.

Taboada EN, Ross SL, Mutschall SK, Mackinnon JM, Roberts MJ, Buchanan CJ, Kruczkiewicz P, Jokinen CC, Thomas JE, Nash JH, Gannon VP, Marshall B, Pollari

F, Clark CG. Development and validation of a comparative genomic fingerprinting method for high-resolution genotyping of *Campylobacter jejuni*. J Clin Microbiol. 2012 Mar;50(3):788-97.

Terra VS, Mills DC, Yates LE, Abouelhadid S, Cuccui J, Wren BW. Recent developments in bacterial protein glycan coupling technology and glycoconjugate vaccine design. J Med Microbiol. 2012 Apr 19. [Epub ahead of print]

Teunis P, van Eijkeren J, Ang C, van Duynhoven Y, Simonsen J, Strid M, van Pelt W. Biomarker dynamics: estimating infection rates from serological data. Stat Med. 2012 Mar 15. doi: 10.1002/sim.5322. [Epub ahead of print]

Ugarte-Ruiz M, Gómez-Barrero S, Porrero M, Alvarez J, García M, Comerón M, Wassenaar T, Domínguez L. Evaluation of four protocols for the detection and isolation of thermophilic *Campylobacter* from different matrices. J Appl Microbiol. 2012 Apr 25. doi: 10.1111/j.1365-2672.2012.05323.x. [Epub ahead of print]

van Bunnik BA, Katsma WE, Wagenaar JA, Jacobs-Reitsma WF, de Jong MC. Acidification of drinking water inhibits indirect transmission, but not direct transmission of *Campylobacter* between broilers. Prev Vet Med. 2012 Apr 18. [Epub ahead of print]

van Gerwe TJ. Poultry meat as a source of human campylobacteriosis. Tijdschr Diergeneeskd. 2012 Mar 1;137(3):172-6. Review.

Wen X, Dickson JS. Survival of *Campylobacter jejuni* and *Salmonella* enteric Typhimurium in vacuum-packed, moisture-enhanced pork. J Food Prot. 2012 Mar;75(3):576-9.

Zeitouni S, Collin O, Andraud M, Ermel G, Kempf I. Fitness of Macrolide Resistant *Campylobacter coli* and *Campylobacter jejuni*. Microb Drug Resist. 2012 Apr;18(2):101-8.

Clostridium

Bader J, Albin A, Stahl U. Spore-forming bacteria and their utilisation as probiotics. Benef Microbes. 2012 Mar 1;3(1):67-75.

Brown JL, Tran-Dinh N, Chapman B. *Clostridium sporogenes* PA 3679 and Its Uses in the Derivation of Thermal Processing Schedules for Low-Acid Shelf-Stable Foods and as a Research Model for Proteolytic *Clostridium botulinum*. J Food Prot. 2012 Apr;75(4):779-92.

Chon JW, Park JS, Hyeon JY, Park C, Song KY, Hong KW, Hwang IG, Kwak HS, Seo KH. Development of Real-Time PCR for the Detection of *Clostridium perfringens* in Meats and Vegetables. J Microbiol Biotechnol. 2012 Apr;22(4):530-4.

Curry SR, Marsh JW, Schlackman JL, Harrison LH. *Clostridium difficile* prevalence in uncooked ground meat products from Pittsburgh, PA. *Appl Environ Microbiol*. 2012 Apr 13. [Epub ahead of print]

Hensgens MP, Keessen EC, Squire MM, Riley TV, Koene MG, de Boer E, Lipman LJ, Kuijper EJ; on behalf of European Society of Clinical Microbiology and Infectious Diseases Study Group for *Clostridium difficile* (ESGCD). *Clostridium difficile* infection in the community: a zoonotic disease? *Clin Microbiol Infect*. 2012 Mar 23. doi: 10.1111/j.1469-0691.2012.03853.x. [Epub ahead of print]

Miyamoto K, Li J, McClane BA. Enterotoxigenic *Clostridium perfringens*: Detection and Identification. *Microbes Environ*. 2012 Apr 14. [Epub ahead of print]

Márquez-González M, Cabrera-Díaz E, Hardin MD, Harris KB, Lucia LM, Castillo A. Survival and Germination of *Clostridium perfringens* Spores during Heating and Cooling of Ground Pork. *J Food Prot*. 2012 Apr;75(4):682-9.

Sevenier V, Delannoy S, André S, Fach P, Remize F. Prevalence of *Clostridium botulinum* and thermophilic heat-resistant spores in raw carrots and green beans used in French canning industry. *Int J Food Microbiol*. 2012 Apr 16;155(3):263-8.

Stabler RA, Dawson LF, Valiente E, Cairns MD, Martin MJ, Donahue EH, Riley TV, Songer JG, Kuijper EJ, Dingle KE, Wren BW. Macro and micro diversity of *Clostridium difficile* isolates from diverse sources and geographical locations. *PLoS One*. 2012;7(3):e31559.

Listeria monocytogenes

Ait-Ouazzou A, Mañas P, Condón S, Pagán R, García-Gonzalo D. Role of general stress-response alternative sigma factors $\sigma(S)$ (RpoS) and $\sigma(B)$ (SigB) in bacterial heat resistance as a function of treatment medium pH. *Int J Food Microbiol*. 2012 Feb 15;153(3):358-64.

Augustin JC, Czarnecka-Kwasiborski A. Single-Cell Growth Probability of *Listeria monocytogenes* at Suboptimal Temperature, pH, and Water Activity. *Front Microbiol*. 2012;3:157. Epub 2012 Apr 24.

Bae YM, Baek SY, Lee SY. Resistance of pathogenic bacteria on the surface of stainless steel depending on attachment form and efficacy of chemical sanitizers. *Int J Food Microbiol*. 2012 Feb 15;153(3):465-73.

Barbosa J, Borges S, Magalhães R, Ferreira V, Santos I, Silva J, Almeida G, Gibbs P, Teixeira P. Behaviour of *Listeria monocytogenes* isolates through gastro-intestinal tract passage simulation, before and after two sub-lethal stresses. *Food Microbiol*. 2012 May;30(1):24-8.

Bergholz TM, Bowen B, Wiedmann M, Boor KJ. *Listeria monocytogenes* Shows Temperature-Dependent and -Independent Responses to Salt Stress, Including

Responses That Induce Cross-Protection against Other Stresses. *Appl Environ Microbiol.* 2012 Apr;78(8):2602-12.

Commeau N, Cornu M, Albert I, Denis JB, Parent E. Hierarchical Bayesian models to assess between- and within-batch variability of pathogen contamination in food. *Risk Anal.* 2012 Mar;32(3):395-415.

Fox EM, deLappe N, Garvey P, McKeown P, Cormican M, Leonard N, Jordan K. PFGE analysis of *Listeria monocytogenes* isolates of clinical, animal, food and environmental origin from Ireland. *J Med Microbiol.* 2012 Apr;61(Pt 4):540-7.

Greenhow TL, Hung YY, Herz AM. Changing epidemiology of bacteremia in infants aged 1 week to 3 months. *Pediatrics.* 2012 Mar;129(3):e590-6.

Hospital XF, Hierro E, Fernández M. Survival of *Listeria innocua* in dry fermented sausages and changes in the typical microbiota and volatile profile as affected by the concentration of nitrate and nitrite. *Int J Food Microbiol.* 2012 Feb 15;153(3):395-401.

Ivy RA, Wiedmann M, Boor KJ. *Listeria monocytogenes* Grown at 7°C Shows Reduced Acid Survival and an Altered Transcriptional Response to Acid Shock Compared to *L. monocytogenes* grown at 37°C. *Appl Environ Microbiol.* 2012 Mar 23. [Epub ahead of print]

Jadhav S, Bhave M, Palombo EA. Methods used for the detection and subtyping of *Listeria monocytogenes*. *J Microbiol Methods.* 2012 Mar;88(3):327-41.

Knabel SJ, Reimer A, Verghese B, Lok M, Ziegler J, Farber J, Pagotto F, Graham M, Nadon CA; the Canadian Public Health Laboratory Network (CPHLN), Gilmour MW. Sequence Typing Confirms that a Predominant *Listeria monocytogenes* Clone Caused Human Listeriosis Cases and Outbreaks in Canada from 1988 to 2010. *J Clin Microbiol.* 2012 May;50(5):1748-1751.

Laksanalamai P, Jackson SA, Mammel MK, Datta AR. High Density Microarray Analysis Reveals New Insights into Genetic Footprints of *Listeria monocytogenes* Strains Involved in Listeriosis Outbreaks. *PLoS One.* 2012;7(3):e32896. Epub 2012 Mar 21.

Pezdiric KB, Hure AJ, Blumfield ML, Collins CE. *Listeria monocytogenes* and diet during pregnancy; balancing nutrient intake adequacy v. adverse pregnancy outcomes. *Public Health Nutr.* 2012 Mar 8:1-8.

Pérez-Rodríguez F, Zwietering MH. Application of the Central Limit Theorem in microbial risk assessment: high number of servings reduces the Coefficient of Variation of food-borne burden-of-illness. *Int J Food Microbiol.* 2012 Feb 15;153(3):413-9.

Sant'Ana AS, Franco BD, Schaffner DW. Modeling the growth rate and lag time of different strains of *Salmonella enterica* and *Listeria monocytogenes* in ready-to-eat lettuce. *Food Microbiol.* 2012 May;30(1):267-73.

Sauders BD, Overdeest J, Fortes E, Windham K, Schukken Y, Lembo A, Wiedmann M. Diversity of *Listeria* species in urban and natural environments. *Appl Environ Microbiol*. 2012 Apr 13. [Epub ahead of print]

Tortajada C, Porta R, Riba M, Santoma MJ, Palacín E, Español M. [Nosocomial outbreak due to *Listeria monocytogenes* in a neonatal unit]. *Enferm Infecc Microbiol Clin*. 2012 Mar;30(3):143-6. Epub 2012 Jan 28. Spanish.

Wieczorek K, Dmowska K, Osek J. Prevalence, characterization, and antimicrobial resistance of *Listeria monocytogenes* isolates from bovine hides and carcasses. *Appl Environ Microbiol*. 2012 Mar;78(6):2043-5.

Zhu L, Stewart D, Reineke K, Ravishankar S, Palumbo S, Cirigliano M, Tortorello M. Comparison of swab transport media for recovery of *Listeria monocytogenes* from environmental samples. *J Food Prot*. 2012 Mar;75(3):580-4.

Norovirus

Brassard J, Gagné MJ, Génereux M, Côté C. Detection of human food-borne and zoonotic viruses on irrigated, field-grown strawberries. *Appl Environ Microbiol*. 2012 May;78(10):3763-6.

Bull RA, Eden JS, Luciani F, McElroy K, Rawlinson WD, White PA. Contribution of intra- and interhost dynamics to norovirus evolution. *J Virol*. 2012 Mar;86(6):3219-29.

Centers for Disease Control and Prevention (CDC). Notes from the field: norovirus infections associated with frozen raw oysters - Washington, 2011. *MMWR Morb Mortal Wkly Rep*. 2012 Feb 17;61(6):110.

Dancho BA, Chen H, Kingsley DH. Discrimination between infectious and non-infectious human norovirus using porcine gastric mucin. *Int J Food Microbiol*. 2012 Apr 16;155(3):222-6.

Desai R, Hembree CD, Handel A, Matthews JE, Dickey BW, McDonald S, Hall AJ, Parashar UD, Leon JS, Lopman B. Severe outcomes are associated with genogroup 2 genotype 4 norovirus outbreaks: A systematic literature review. *Clin Infect Dis*. 2012 Apr 4. [Epub ahead of print]

Flannery J, Keaveney S, Rajko-Nenow P, O'Flaherty V, Doré W. Concentration of Norovirus during Wastewater Treatment and Its Impact on Oyster Contamination. *Appl Environ Microbiol*. 2012 May;78(9):3400-6.

Gerhardt A, Hammer TR, Balluff C, Mucha H, Hofer D. A model of the transmission of micro-organisms in a public setting and its correlation to pathogen infection risks. *J Appl Microbiol*. 2012 Mar;112(3):614-21.

Gibson KE, Crandall PG, Ricke SC. Removal and transfer of viruses on food contact surfaces by cleaning cloths. *Appl Environ Microbiol*. 2012 May;78(9):3037-44.

Haramoto E, Katayama H, Asami M, Akiba M. Development of a novel method for simultaneous concentration of viruses and protozoa from a single water sample. *J Virol Methods*. 2012 Jun;182(1-2):62-9.

Heijne JC, Rondy M, Verhoef L, Wallinga J, Kretzschmar M, Low N, Koopmans M, Teunis PF. Quantifying transmission of norovirus during an outbreak. *Epidemiology*. 2012 Mar;23(2):277-84.

Hoffmann D, Hutzenthaler M, Seebach J, Panning M, Umgelter A, Menzel H, Protzer U, Metzler D. Norovirus GII.4 and GII.7 capsid sequences undergo positive selection in chronically infected patients. *Infect Genet Evol*. 2012 Mar;12(2):461-6.

Kirby A, Iturriza-Gómara M. Norovirus diagnostics: options, applications and interpretations. *Expert Rev Anti Infect Ther*. 2012 Apr;10(4):423-33.

Li D, Baert L, Xia M, Zhong W, Van Coillie E, Jiang X, Uyttendaele M. Evaluation of methods measuring the capsid integrity and/or functions of noroviruses by heat inactivation. *J Virol Methods*. 2012 Apr;181(1):1-5.

Longtin Y, Voss A, Allegranzi B, Pittet D. Norovirus outbreaks and alcohol-based handrub solutions: association does not prove causation. *Am J Infect Control*. 2012 Mar;40(2):191; author reply 192.

Matthews JE, Dickey BW, Miller RD, Felzer JR, Dawson BP, Lee AS, Rocks JJ, Kiel J, Montes JS, Moe CL, Eisenberg JN, Leon JS. The epidemiology of published norovirus outbreaks: a review of risk factors associated with attack rate and genogroup. *Epidemiol Infect*. 2012 Mar 26:1-12.

Maunula L, Söderberg K, Vahtera H, Vuorilehto VP, von Bonsdorff CH, Valtari M, Laakso T, Lahti K. Presence of human noro- and adenoviruses in river and treated wastewater, a longitudinal study and method comparison. *J Water Health*. 2012 Mar;10(1):87-99.

McAllister G, Holmes A, Garcia L, Cameron F, Cloy K, Danial J, Cepeda JA, Simmonds P, Templeton KE. Molecular epidemiology of norovirus in Edinburgh healthcare facilities, Scotland 2007-2011. *Epidemiol Infect*. 2012 Feb 6:1-9.

Nenonen NP, Hannoun C, Larsson CU, Bergström T. Marked genomic diversity of norovirus genogroup I strains in a waterborne outbreak. *Appl Environ Microbiol*. 2012 Mar;78(6):1846-52.

Pinto P, Wang Q, Chen N, Dubovi EJ, Daniels JB, Millward LM, Buonavoglia C, Martella V, Saif LJ. Discovery and genomic characterization of noroviruses from a gastroenteritis outbreak in domestic cats in the US. *PLoS One*. 2012;7(2):e32739.

Richards GP. Critical Review of Norovirus Surrogates in Food Safety Research: Rationale for Considering Volunteer Studies. *Food Environ Virol*. 2012 Mar;4(1):6-13.

Seo K, Lee JE, Lim MY, Ko G. Effect of temperature, pH, and NaCl on the inactivation kinetics of murine norovirus. *J Food Prot.* 2012 Mar;75(3):533-40.

Tuladhar E, Bouwknecht M, Zwietering MH, Koopmans M, Duizer E. Thermal stability of structurally different viruses with proven or potential relevance to food safety. *J Appl Microbiol.* 2012 May;112(5):1050-1057.

Salmonella

Bae YM, Baek SY, Lee SY. Resistance of pathogenic bacteria on the surface of stainless steel depending on attachment form and efficacy of chemical sanitizers. *Int J Food Microbiol.* 2012 Feb 15;153(3):465-73.

Berge AC, Wierup M. Nutritional strategies to combat *Salmonella* in mono-gastric food animal production. *Animal.* 2012 Apr;6(4):557-64.

Betancor L, Yim L, Martínez A, Fookes M, Sasias S, Schelotto F, Thomson N, Maskell D, Chabalgoity JA. Genomic Comparison of the Closely Related *Salmonella enterica* Serovars Enteritidis and Dublin. *Open Microbiol J.* 2012;6:5-13..

Bolton DJ, O'Neill CJ, Fanning S. A Preliminary Study of *Salmonella*, Verocytotoxigenic *Escherichia coli*/*Escherichia coli* O157 and *Campylobacter* on Four Mixed Farms. *Zoonoses Public Health.* 2012 May;59(3):217-28.

Brashears MM, Garmyn AJ, Brooks JC, Harris D, Loneragan G, Echeverry A, Jackson TE, Mehaffey JM, Miller MF. Microbial quality of condensation in fresh and ready-to-eat processing facilities. *Meat Sci.* 2012 Mar;90(3):728-32.

Cevallos-Cevallos JM, Danyluk MD, Gu G, Vallad GE, van Bruggen AH. Dispersal of *Salmonella* Typhimurium by rain splash onto tomato plants. *J Food Prot.* 2012 Mar;75(3):472-9.

Condell O, Iversen C, Cooney S, Power KA, Walsh C, Burgess C, Fanning S. Efficacy of Biocides Used in the Modern Food Industry To Control *Salmonella enterica*, and Links between Biocide Tolerance and Resistance to Clinically Relevant Antimicrobial Compounds. *Appl Environ Microbiol.* 2012 May;78(9):3087-97.

David R. Bacterial pathogenesis: A competitive edge for *Salmonella*. *Nat Rev Microbiol.* 2012 Apr 11;10(5):309.

de Jong A, Thomas V, Simjee S, Godinho K, Schiessl B, Klein U, Butty P, Vallé M, Marion H, Shryock TR. Pan-European monitoring of susceptibility to human-use antimicrobial agents in enteric bacteria isolated from healthy food-producing animals. *J Antimicrob Chemother.* 2012 Mar;67(3):638-51.

Deng X, Li Z, Zhang W. Transcriptome sequencing of *Salmonella enteric* serovar Enteritidis under desiccation and starvation stress in peanut oil. *Food Microbiol.* 2012 May;30(1):311-5.

Gajraj R, Pooransingh S, Hawker JI, Olowokure B. Multiple outbreaks of *Salmonella braenderup* associated with consumption of iceberg lettuce. *Int J Environ Health Res.* 2012 Apr;22(2):150-5.

Gossner CM, van Cauteren D, Le Hello S, Weill FX, Terrien E, Tessier S, Janin C, Brisabois A, Dusch V, Vaillant V, Jourdan-da Silva N. Nationwide outbreak of *Salmonella enterica* serotype 4,[5],12:i:- infection associated with consumption of dried pork sausage, France, November to December 2011. *Euro Surveill.* 2012 Feb 2;17(5). pii: 20071. Erratum in: *Euro Surveill.* 2012;17(6). pii: 20078. Vaillant, V [added]. PubMed PMID: 22321138.

Hauser E, Tietze E, Helmuth R, Junker E, Prager R, Schroeter A, Rabsch W, Fruth A, Toboldt A, Malorny B. Clonal dissemination of *Salmonella enterica* serovar Infantis in Germany. *Foodborne Pathog Dis.* 2012 Apr;9(4):352-60.

Havelaar AH, Haagsma JA, Mangen MJ, Kemmeren JM, Verhoef LP, Vijgen SM, Wilson M, Friesema IH, Kortbeek LM, van Duynhoven YT, van Pelt W. Disease burden of foodborne pathogens in the Netherlands, 2009. *Int J Food Microbiol.* 2012 Apr 5. [Epub ahead of print]

Ivanek R, Osterberg J, Gautam R, Sternberg Lewerin S. *Salmonella* Fecal Shedding and Immune Responses are Dose- and Serotype- Dependent in Pigs. *PLoS One.* 2012;7(4):e34660. Epub 2012 Apr 16.

Jeong S, Marks BP, Ryser ET, Harte JB. The effect of X-ray irradiation on *Salmonella* inactivation and sensory quality of almonds and walnuts as a function of water activity. *Int J Food Microbiol.* 2012 Feb 15;153(3):365-71.

Jones TF, Gerner-Smidt P. Nonculture diagnostic tests for enteric diseases. *Emerg Infect Dis.* 2012 Mar;18(3):513-4.

Kisluk G, Yaron S. Presence and persistence of *Salmonella enterica* serotype Typhimurium on the phyllosphere and rhizosphere of spray irrigated parsley. *Appl Environ Microbiol.* 2012 Mar 23. [Epub ahead of print]

Kröger C, Dillon SC, Cameron AD, Papenfort K, Sivasankaran SK, Hokamp K, Chao Y, Sittka A, Hébrard M, Händler K, Colgan A, Leekitcharoenphon P, Langridge GC, Lohan AJ, Loftus B, Lucchini S, Ussery DW, Dorman CJ, Thomson NR, Vogel J, Hinton JC. The transcriptional landscape and small RNAs of *Salmonella enterica* serovar Typhimurium. *Proc Natl Acad Sci U S A.* 2012 Apr 25. [Epub ahead of print]

Leekitcharoenphon P, Lukjancenko O, Friis C, Aarestrup FM, Ussery DW. Genomic variation in *Salmonella enterica* core genes for epidemiological typing. *BMC Genomics.* 2012 Mar 12;13(1):88.

Lungu B, Waltman WD, Berghaus RD, Hofacre CL. Comparison of a Real-Time PCR Method with a Culture Method for the Detection of *Salmonella enterica* Serotype Enteritidis in Naturally Contaminated Environmental Samples from Integrated Poultry Houses. *J Food Prot.* 2012 Apr;75(4):743-7.

Mather AE, Matthews L, Mellor DJ, Reeve R, Denwood MJ, Boerlin P, Reid-Smith RJ, Brown DJ, Coia JE, Browning LM, Haydon DT, Reid SW. An ecological approach to assessing the epidemiology of antimicrobial resistance in animal and human populations. *Proc Biol Sci.* 2012 Apr 22;279(1733):1630-9.

Moreira RG, Puerta-Gomez AF, Kim J, Castell-Perez ME. Factors Affecting Radiation D-Values (D(10)) of an *Escherichia Coli* Cocktail and *Salmonella* Typhimurium LT2 Inoculated in Fresh Produce. *J Food Sci.* 2012 Apr;77(4):E104-11. doi: 10.1111/j.1750-3841.2011.02603.x.

Morey A, Singh M. Low-temperature survival of *Salmonella* spp. in a model food system with natural microflora. *Foodborne Pathog Dis.* 2012 Mar;9(3):218-23.

Méroc E, Strubbe M, Vangroenweghe F, Czaplicki G, Vermeersch K, Hooyberghs J, Van der Stede Y. Evaluation of the *Salmonella* surveillance program in Belgian pig farms. *Prev Vet Med.* 2012 Apr 7. [Epub ahead of print]

Müller K, Aabo S, Birk T, Mordhorst H, Bjarnadóttir B, Agersø Y. Survival and growth of epidemically successful and nonsuccessful *Salmonella enterica* clones after freezing and dehydration. *J Food Prot.* 2012 Mar;75(3):456-64.

Najjar Z, Furlong C, Stephens N, Shadbolt C, Maywood P, Conaty S, Hogg G. An outbreak of *Salmonella* Infantis gastroenteritis in a residential aged care facility associated with thickened fluids. *Epidemiol Infect.* 2012 Feb 16:1-9. [Epub ahead of print]

Neetoo H, Chen H. High pressure inactivation of *Salmonella* on Jalapeño and Serrano peppers destined for direct consumption or as ingredients in Mexican salsa and guacamole. *Int J Food Microbiol.* 2012 Mar 26. [Epub ahead of print].

Neetoo H, Lu Y, Wu C, Chen H. Use of high hydrostatic pressure to inactivate *Escherichia coli* O157:H7 and *Salmonella enterica* internalized within and adhered to preharvest contaminated green onions. *Appl Environ Microbiol.* 2012 Mar;78(6):2063-5.

Pao S, Long W 3rd, Kim C, Rafie AR. *Salmonella* population rebound and its prevention on spray washed and non-washed jalapeño peppers and roma tomatoes in humid storage. *Foodborne Pathog Dis.* 2012 Apr;9(4):361-6.

Papadopoulou OS, Chorianopoulos NG, Gkana EN, Grounta AV, Koutsoumanis KP, Nychas GJ. Transfer of foodborne pathogenic bacteria to non-inoculated beef fillets through meat mincing machine. *Meat Sci.* 2012 Mar;90(3):865-9.

Quigley L, O'Sullivan O, Beresford TP, Paul Ross R, Fitzgerald GF, Cotter PD. A comparison of methods used to extract bacterial DNA from raw milk and raw milk cheese. *J Appl Microbiol.* 2012 Mar 27. doi: 10.1111/j.1365-2672.2012.05294.x. [Epub ahead of print].

Zhu J, Li J, Chen J. Survival of Salmonella in home-style mayonnaise and acid solutions as affected by acidulant type and preservatives. J Food Prot. 2012 Mar;75(3):465-71.

Álvarez-Fernández E, Alonso-Calleja C, García-Fernández C, Capita R. Prevalence and antimicrobial resistance of *Salmonella* serotypes isolated from poultry in Spain: comparison between 1993 and 2006. Int J Food Microbiol. 2012 Feb 15;153(3):281-7.

Staphylococcus aureus

Argudín MA, Mendoza MC, González-Hevia MA, Bances M, Guerra B, Rodicio MR. Genotypes, Exotoxin Gene Content, and Antimicrobial Resistance of *Staphylococcus aureus* Strains Recovered from Foods and Food Handlers. Appl Environ Microbiol. 2012 Apr;78(8):2930-5.

Jouy E, Le Roux A, Kéranflec'h A, Granier SA, Laurent F, Kempf I, Brisabois A, Cariolet R, Chauvin C. Methicillin-resistant *Staphylococcus aureus* ST398 contamination and transmission in pigs after a low dose inoculation. Lett Appl Microbiol. 2012 Mar 10. doi: 10.1111/j.1472-765X.2012.03239.x.

Maisch T, Spannberger F, Regensburger J, Felgenträger A, Bäumler W. Fast and effective: intense pulse light photodynamic inactivation of bacteria. J Ind Microbiol Biotechnol. 2012 Feb 22. [Epub ahead of print]

Morcillo A, Castro B, Rodríguez-Álvarez C, González JC, Sierra A, Montesinos MI, Abreu R, Arias A. Prevalence and characteristics of methicillin-resistant *Staphylococcus aureus* in pigs and pig workers in Tenerife, Spain. Foodborne Pathog Dis. 2012 Mar;9(3):207-10.

Pantosti A. Methicillin-Resistant *Staphylococcus aureus* Associated with Animals and Its Relevance to Human Health. Front Microbiol. 2012;3:127.

Porrero MC, Wassenaar TM, Gómez-Barrero S, García M, Bárcena C, Alvarez J, Sáez-Llorente JL, Fernández-Garayzabal JF, Moreno MA, Domínguez L. Detection of methicillin-resistant *Staphylococcus aureus* in Iberian pigs. Lett Appl Microbiol. 2012 Apr;54(4):280-5. doi: 10.1111/j.1472-765X.2012.03207.x. Epub 2012 Feb 13. PubMed PMID: 22251394.

Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, Pearson T, Waters AE, Foster JT, Schupp J, Gillece J, Driebe E, Liu CM, Springer B, Zdovc I, Battisti A, Franco A, Zmudzki J, Schwarz S, Butaye P, Jouy E, Pomba C, Porrero MC, Ruimy R, Smith TC, Robinson DA, Weese JS, Arriola CS, Yu F, Laurent F, Keim P, Skov R, Aarestrup FM. *Staphylococcus aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. MBio. 2012 Feb 21;3(1). pii:e00305-11. doi: 10.1128/mBio.00305-11.

Shahraz F, Dadkhah H, Khaksar R, Mahmoudzadeh M, Hosseini H, Kamran M, Bourke P. Analysis of antibiotic resistance patterns and detection of *mecA* gene in

Staphylococcus aureus isolated from packaged hamburger. Meat Sci. 2012 Mar;90(3):759-63.

Weese JS, Hannon SJ, Booker CW, Gow S, Avery BP, Reid-Smith RJ. The prevalence of methicillin-resistant *Staphylococcus aureus* colonization in feedlot cattle. Zoonoses Public Health. 2012 Mar;59(2):144-7.

Toxoplasma

Arias I, Sorlozano A, Villegas E, de Dios Luna J, McKenney K, Cervilla J, Gutierrez B, Gutierrez J. Infectious agents associated with schizophrenia: a meta-analysis. Schizophr Res. 2012 Apr;136(1-3):128-36.

Bayarri S, Gracia MJ, Pérez-Arquillué C, Lázaro R, Herrera A. *Toxoplasma gondii* in commercially available pork meat and cured ham: a contribution to risk assessment for consumers. J Food Prot. 2012 Mar;75(3):597-600.

Dubey JP, Hill DE, Rozeboom DW, Rajendran C, Choudhary S, Ferreira LR, Kwok OC, Su C. High prevalence and genotypes of *Toxoplasma gondii* isolated from organic pigs in northern USA. Vet Parasitol. 2012 Mar 15. [Epub ahead of print]

Havelaar AH, Haagsma JA, Mangen MJ, Kemmeren JM, Verhoef LP, Vijgen SM, Wilson M, Friesema IH, Kortbeek LM, van Duynhoven YT, van Pelt W. Disease burden of foodborne pathogens in the Netherlands, 2009. Int J Food Microbiol. 2012 Apr 5. [Epub ahead of print]

Opsteegh M, Haveman R, Swart AN, Mensink-Beerepoot ME, Hofhuis A, Langelaar MF, van der Giessen JW. Seroprevalence and risk factors for *Toxoplasma gondii* infection in domestic cats in The Netherlands. Prev Vet Med. 2012 May 1;104(3-4):317-26. Epub 2012 Feb 2. PubMed PMID: 22305876.

Petersen E, Kijlstra A, Stanford M. Epidemiology of ocular toxoplasmosis. Ocul Immunol Inflamm. 2012 Apr;20(2):68-75.

Robert-Gangneux F, Dardé ML. Epidemiology of and diagnostic strategies for toxoplasmosis. Clin Microbiol Rev. 2012 Apr;25(2):264-96.

Sakikawa M, Noda S, Hanaoka M, Nakayama H, Hojo S, Kakinoki S, Nakata M, Yasuda T, Ikenoue T, Kojima T. Anti-Toxoplasma antibody prevalence, primary infection rate, and risk factors in a study of toxoplasmosis in 4,466 pregnant women in Japan. Clin Vaccine Immunol. 2012 Mar;19(3):365-7.

Sullivan WJ Jr, Jeffers V. Mechanisms of *Toxoplasma gondii* persistence and latency. FEMS Microbiol Rev. 2012 May;36(3):717-33.

Useo R, Husson F, De Coninck J, Khaldi S, Gervais P. A New Alternative In Vitro Method for Quantification of *Toxoplasma gondii* Infectivity. J Parasitol. 2012 Apr;98(2):299-303.

Villena I, Durand B, Aubert D, Blaga R, Geers R, Thomas M, Perret C, Alliot A, Escotte-Binet S, Thébault A, Boireau P, Halos L. New strategy for the survey of *Toxoplasma gondii* in meat for human consumption. *Vet Parasitol.* 2012 Feb 10;183(3-4):203-8.

VTEC/STEC

Anklam KS, Kanankege KS, Gonzales TK, Kaspar CW, Döpfer D. Rapid and Reliable Detection of Shiga Toxin-Producing *Escherichia coli* by Real-Time Multiplex PCR. *J Food Prot.* 2012 Apr;75(4):643-50.

Asakura H, Saito E, Momose Y, Ekawa T, Sawada M, Yamamoto A, Hasegawa A, Iwahori J, Tsutsui T, Osaka K, Matsushita T, Kakinuma M, Motoyama K, Hayama Y, Kitamoto H, Igimi S, Kasuga F. Prevalence and growth kinetics of Shiga toxin-producing *Escherichia coli* (STEC) in bovine offal products in Japan. *Epidemiol Infect.* 2012 Apr;140(4):655-64.

Auvray F, Dilasser F, Bibbal D, Kérourédan M, Oswald E, Brugère H. French cattle is not a reservoir of the highly virulent enteroaggregative Shiga toxin-producing *Escherichia coli* of serotype O104:H4. *Vet Microbiol.* 2012 Feb 28. [Epub ahead of print]

Bezanson G, Delaquis P, Bach S, McKellar R, Topp E, Gill A, Blais B, Gilmour M. Comparative examination of *Escherichia coli* O157:H7 survival on romaine lettuce and in soil at two independent experimental sites. *J Food Prot.* 2012 Mar;75(3):480-7.

Biliński P, Kapka-Skrzypczak L, Posobkiewicz M, Bondaryk M, Hołownia P, Wojtyła A. Public health hazards in Poland posed by foodstuffs contaminated with *E.Coli* O104:H4 bacterium from the recent European outbreak. *Ann Agric Environ Med.* 2012 Mar 23;19(1):3-10.

Castro-Rosas J, Cerna-Cortés JF, Méndez-Reyes E, Lopez-Hernandez D, Gómez-Aldapa CA, Estrada-Garcia T. Presence of faecal coliforms, *Escherichia coli* and diarrheagenic *E. coli* pathotypes in ready-to-eat salads, from an area where crops are irrigated with untreated sewage water. *Int J Food Microbiol.* 2012 Mar 30. [Epub ahead of print]

Cross P, Rigby D, Edwards-Jones G. Eliciting expert opinion on the effectiveness and practicality of interventions in the farm and rural environment to reduce human exposure to *Escherichia coli* O157. *Epidemiol Infect.* 2012 Apr;140(4):643-54.

Fernandez-Brando RJ, Miliwebsky E, Mejías MP, Baschkier A, Panek CA, Abrey-Recalde MJ, Cabrera G, Ramos MV, Rivas M, Palermo MS. Shiga toxin-producing *Escherichia coli* O157:H7 show an increased pathogenicity in mice after the passage through the gastrointestinal tract of the same host. *J Med Microbiol.* 2012 Mar 15. [Epub ahead of print]

Harris D, Brashears MM, Garmyn AJ, Brooks JC, Miller MF. Microbiological and organoleptic characteristics of beef trim and ground beef treated with acetic acid, lactic acid, acidified sodium chlorite, or sterile water in a simulated commercial processing environment to reduce *Escherichia coli* O157:H7 and *Salmonella*. *Meat Sci.* 2012 Mar;90(3):783-8.

Hegde NV, Jayarao BM, Debroy C. Rapid Detection of the Top Six Non-O157 Shiga Toxin-Producing *Escherichia coli* O Groups in Ground Beef by Flow Cytometry. *J Clin Microbiol.* 2012 Apr 4. [Epub ahead of print]

Hofer E, Stephan R, Reist M, Zweifel C. Application of a Real-Time PCR-Based System for Monitoring of O26, O103, O111, O145 and O157 Shiga Toxin-Producing *Escherichia coli* in Cattle at Slaughter. *Zoonoses Public Health.* 2012 Feb 21. doi: 10.1111/j.1863-2378.2012.01468.x. [Epub ahead of print].

Huang L, Tu SI, Phillips J, Fratamico P. Mathematical Modeling of Growth of Non-O157 Shiga Toxin-Producing *Escherichia coli* in Raw Ground Beef. *J Food Sci.* 2012 Apr;77(4):M217-25.

Januszkiewicz A, Szych J, Rastawicki W, Wołkowicz T, Chróst A, Leszczyńska B, Kuźma E, Roszkowska-Blaim M, Gierczyński R. Molecular epidemiology of a household outbreak of Shiga-toxin-producing *Escherichia coli* in Poland due to secondary transmission of STEC O104:H4 from Germany. *J Med Microbiol.* 2012 Apr;61(Pt 4):552-8.

Jenkins C, Lawson A, Cheasty T, Bolton E, Smith G. Assessment of a real-time PCR for the detection and characterisation of Verocytotoxigenic *Escherichia coli*. *J Med Microbiol.* 2012 Apr 19. [Epub ahead of print]

Jinneman KC, Waite-Cusic JG, Yoshitomi KJ. Evaluation of shiga toxin-producing *Escherichia coli* (STEC) method for the detection and identification of STEC O104 strains from sprouts. *Food Microbiol.* 2012 May;30(1):321-8.

King LA, Nogareda F, Weill FX, Mariani-Kurkdjian P, Loukiadis E, Gault G, Jourdan-Dasilva N, Bingen E, Macé M, Thevenot D, Ong N, Castor C, Noël H, Van Cauteren D, Charron M, Vaillant V, Aldabe B, Goulet V, Delmas G, Couturier E, Le Strat Y, Combe C, Delmas Y, Terrier F, Vendrely B, Rolland P, de Valk H. Outbreak of Shiga Toxin-Producing *Escherichia coli* O104:H4 Associated With Organic Fenugreek Sprouts, France, June 2011. *Clin Infect Dis.* 2012 Mar 28. [Epub ahead of print]

Lal A, Hales S, French N, Baker MG. Seasonality in human zoonotic enteric diseases: a systematic review. *PLoS One.* 2012;7(4):e31883.

Lee K, French NP, Jones G, Hara-Kudo Y, Iyoda S, Kobayashi H, Sugita-Konishi Y, Tsubone H, Kumagai S. Variation in Stress Resistance Patterns among stx Genotypes and Genetic Lineages of Shiga Toxin-Producing *Escherichia coli* O157. *Appl Environ Microbiol.* 2012 May;78(9):3361-8.

Mora A, López C, Dhahi G, López-Beceiro AM, Fidalgo LE, Díaz EA, Martínez-Carrasco C, Mamani R, Herrera A, Blanco JE, Blanco M, Blanco J. Seropathotypes, Phylogroups, Stx Subtypes, and Intimin Types of Wildlife-Carried, Shiga Toxin-Producing *Escherichia coli* Strains with the Same Characteristics as Human-Pathogenic Isolates. *Appl Environ Microbiol.* 2012 Apr;78(8):2578-85.

Muniesa M, Hammerl JA, Hertwig S, Appel B, Brüssow H. Shiga toxin-producing *Escherichia coli* O104:H4 - a new challenge for microbiology. *Appl Environ Microbiol.* 2012 Apr 13. [Epub ahead of print]

Neil KP, Biggerstaff G, MacDonald JK, Trees E, Medus C, Musser KA, Stroika SG, Zink D, Sotir MJ. A novel vehicle for transmission of *Escherichia coli* O157:H7 to humans: multistate outbreak of *E. coli* O157:H7 infections associated with consumption of ready-to-bake commercial prepackaged cookie dough—United States, 2009. *Clin Infect Dis.* 2012 Feb 15;54(4):511-8.

Niu YD, Stanford K, Kropinski AM, Ackermann HW, Johnson RP, She YM, Ahmed R, Villegas A, McAllister TA. Genomic, Proteomic and Physiological Characterization of a T5-like Bacteriophage for Control of Shiga Toxin-Producing *Escherichia coli* O157:H7. *PLoS One.* 2012;7(4):e34585.

Rode TM, Holck A, Axelsson L, Høy M, Heir E. Shiga toxigenic *Escherichia coli* show strain dependent reductions under dry-fermented sausage production and post-processing conditions. *Int J Food Microbiol.* 2012 Apr 16;155(3):227-33.

Staples M, Graham RM, Doyle CJ, Smith HV, Jennison AV. Prolonged and mixed non-O157 *Escherichia coli* infection in an Australian household. *Clin Microbiol Infect.* 2012 May;18(5):E140-3.

Stokes MO, Cottell JL, Piddock LJ, Wu G, Wootton M, Mevius DJ, Randall LP, Teale CJ, Fielder MD, Coldham NG. Detection and characterization of pCT-like plasmid vectors for blaCTX-M-14 in *Escherichia coli* isolates from humans, turkeys and cattle in England and Wales. *J Antimicrob Chemother.* 2012 Apr 18. [Epub ahead of print]

Thomas KM, McCann MS, Collery MM, Logan A, Whyte P, McDowell DA, Duffy G. Tracking verocytotoxigenic *Escherichia coli* O157, O26, O111, O103 and O145 in Irish cattle. *Int J Food Microbiol.* 2012 Feb 15;153(3):288-96.

Toljander J, Dovärn A, Andersson Y, Ivarsson S, Lindqvist R. Public health burden due to infections by verocytotoxin-producing *Escherichia coli* (VTEC) and *Campylobacter* spp. as estimated by cost of illness and different approaches to model disability-adjusted life years. *Scand J Public Health.* 2012 Feb 6. [Epub ahead of print]

Wang F, Jiang L, Yang Q, Prinyawiwatkul W, Ge B. Rapid and Specific Detection of *Escherichia coli* Serogroups O26, O45, O103, O111, O121, O145, and O157 in Ground Beef, Beef Trim, and Produce by Loop-Mediated Isothermal Amplification. *Appl Environ Microbiol.* 2012 Apr;78(8):2727-36.

Zhou B, Luo Y, Millner P, Feng H. Sanitation and design of lettuce coring knives for minimizing *Escherichia coli* O157:H7 contamination. J Food Prot. 2012 Mar;75(3):563-6.