

**ADVISORY COMMITTEE ON THE MICROBIOLOGICAL SAFETY OF FOOD**

**Items of possible interest from the literature**

A list of items from the literature which may be of interest to Members is attached.

**Secretariat  
March 2008**

## ***Bacillus cereus***

Andrup L, Barfod KK, Jensen GB, Smidt L. Detection of large plasmids from the *Bacillus cereus* group. *Plasmid* 2008 Jan 5 [Epub ahead of print]

Tudela E, Croizé J, Lagier A, Mallaret MR. [Microbiological monitoring of milk samples and surface samples in a hospital infant formula room.] *Pathol Biol (Paris)* 2008 Jan 3 [Epub ahead of print] French.

Auger S, Galleron N, Bidnenko E, Ehrlich SD, Lapidus A, Sorokin A. The genetically remote pathogenic strain NVH391-98 of the *Bacillus cereus* group is representative of a cluster of thermophilic strains. *Appl Environ Microbiol* 2008; **74(4)**: 1276-80.

Baron F, Cochet MF, Grosset N, Madec MN, Briandet R, Dessaigne S, Chevalier S, Gautier M, Jan S. Isolation and characterization of a psychrotolerant toxin producer, *Bacillus weihenstephanensis*, in liquid egg products. *J Food Prot* 2007; **70(12)**: 2782-91.

Yang IC, Shih DY, Wang JY, Pani TM. Development of rapid real-time PCR and most-probable-number real-time PCR assays to quantify enterotoxigenic strains of the species in the *Bacillus cereus* group. *J Food Prot* 2007; **70(12)**: 2774-81.

Cardazzo B, Negrisolo E, Carraro L, Alberghini L, Patarnello T, Giaccone V. Multiple-locus sequence typing and analysis of toxin gene in *Bacillus cereus* food-borne isolates. *Appl Environ Microbiol* 2008; **74(3)**: 850-60.

Kyei-Poku G, Gauthier D, Pang A, van Frankenhuyzen K. Detection of *Bacillus cereus* factors in commercial products of *Bacillus thuringiensis* and expression of diarrheal enterotoxins in a target insect. *Can J Microbiol* 2007; **53(12)**: 1283-90.

Guinebretière MH, Thompson FL, Sorokin A, Normand P, Dawyndt P, Ehling-Schulz M, Svensson B, Sanchis V, Nguyen-The C, Heyndrickx M, De Vos P. Ecological diversification in the *Bacillus cereus* Group. *Environ Microbiol* 2007 Nov 25 [Epub ahead of print].

Al-Khatib MS, Khyami-Horani H, Badran E, Shehabi AA. Incidence and characterization of diarrheal enterotoxins of fecal *Bacillus cereus* isolates associated with diarrhea. *Diagn Microbiol Infect Dis* 2007; **59(4)**: 383-7.

Lapidus A, Goltsman E, Auger S, Galleron N, Ségurens B, Dossat C, Land ML, Broussolle V, Brillard J, Guinebretière MH, Sanchis V, Nguyen-The C, Lereclus D, Richardson P, Wincker P, Weissenbach J, Ehrlich SD, Sorokin A. Extending the *Bacillus cereus* group genomics to putative food-borne pathogens of different toxicity. *Chem Biol Interac* 2008; **171(2)**: 236-49.

## ***Campylobacter***

Elvers KT, Helps CR, Wassenaar TM, Allen VM, Newell DG. Development of a Strain-Specific Molecular Method for Quantitating Individual *Campylobacter* Strains in Mixed Populations. *Appl Environ Microbiol* 2008 Feb 15; [Epub ahead of print]

Wehebrink T, Kemper N, grosse Beilage E, Krieter J. Prevalence of *Campylobacter* spp. and *Yersinia* spp. in the pig production. *Berl Munch Tierarztl Wochenschr* 2008; **121(1-2)**: 27-32.

Verhoeff-Bakkenes L, Arends AP, Snoep JL, Zwietering MH, de Jonge R. Pyruvate relieves the necessity of high induction levels of catalase enables *Campylobacter jejuni* to grow under fully aerobic conditions. *Lett Appl Microbiol* 2008; Feb 7 [Epub ahead of print]

Hald B, Sommer HM, Skovgård H. Use of fly screens to reduce *Campylobacter* spp. Introduction in broiler houses. *Emerg Infect Dis* 2007; **13(12)**: 1951-3.

McDowell SW, Menzies FD, McBride SH, Oza AN, McKenna JP, Gordon AW, Neill SD. *Campylobacter* spp. in conventional broiler flocks in Northern Ireland: Epidemiology and risk factors. *Prev Vet Med* 2008 Feb 2; [Epub ahead of print].

Ghafir Y, China B, Dierick K, De Zutter L, Daube G. Hygiene indicator microorganisms for selected pathogens on beef, pork, and poultry meats in Belgium. *J Food Prot* 2008; **71(1)**: 35-45.

Unicomb LE, O'Reilly LC, Kirk MD, Stafford RJ, Smith HV, Becker NG, Patel MS, Gilbert GL; the Australian *Campylobacter* Subtyping Study Group. Risk factors for infection with *Campylobacter jejuni* flaA genotypes. *Epidemiol Infect* 2008; Jan 21, 1-12 [Epub ahead of print]

Ridley AM, Allen VM, Sharma M, Harris JA, Newell DG. A real time polymerase chain reaction approach for the detection of environmental sources of *Campylobacter* strains colonizing broiler flocks. *Appl Environ Microbiol* 2008 Jan 18 [Epub ahead of print]

Reid AN, Pandey R, Palyada K, Naikare H, Stintzi A. Identification of *Campylobacter jejuni* genes involved in the response to acidic pH and stomach transit. *Appl Environ Microbiol* 2008; Jan [Epub ahead of print]

Van Deun K, Pasmans F, Ducatelle R, Flahou B, Vissenberg K, Martel A, Van den Broeck W, Van Immerseel F, Haesebrouck F. Colonization strategy of *Campylobacter jejuni* results in persistent infection of the chicken gut. *Vet Microbiol* 2007 Dec 3 [Epub ahead of print]

Garénaux A, Jugiau F, Rama F, de Jonge R, Denis M, Federighi M, Ritz M. Survival of *Campylobacter jejuni* Strains from Different Origins Under

Oxidative Stress Conditions: Effect of Temperature. *Curr Microbiol* 2008 Jan 5 [Epub ahead of print]

Guillou S, Leguerinel I, Garrec N, Renard MA, Cappelier JM, Federighi M. Survival of *Campylobacter jejuni* in mineral bottled water according to difference in mineral content: Application of the Weibull model. *Water Res* 2007 Dec 5 [Epub ahead of print]

Peyrat MB, Soumet C, Maris P, Sanders P. Phenotypes and genotypes of *Campylobacter* strains isolated after cleaning and disinfection in poultry slaughterhouses. *Vet Microbiol* 2007 Dec 10; [Epub ahead of print]

Woldermariam E, Bouma A, Vernooij JC, Stegeman A. The sensitivity and specificity of fecal and cecal culture for the detection of *Campylobacter* in Dutch broiler flocks quantified by Bayesian analysis. *Int J Food Microbiol* 2008; **121(3)**: 308-12.

Gormley FJ, Macrae M, Forbes KJ, Ogden ID, Dallas JF, Strachan NJ. Has retail chicken played a role in the decline of human campylobacteriosis? *Appl Environ Microbiol* 2008; **74(2)**: 383-90.

Garrett N, Devane ML, Hudson JA, Nicol C, Ball A, Klena JD, Scholes P, Baker MG, Gilpin BJ, Savill MG. Statistical comparison of *Campylobacter jejuni* subtypes from human cases and environmental sources. *J Appl Microbiol* 2007; **103(6)**: 2113-21.

Poly F, Guerry P. Pathogenesis of *Campylobacter*. *Curr Opin Gastroenterol* 2008; **24(1)**: 27-31. Review.

Lindqvist R, Lindblad M. Quantitative risk assessment of thermophilic *Campylobacter* spp. and cross-contamination during handling of raw broiler chickens evaluating strategies at the producer level to reduce human campylobacteriosis in Sweden. *Int J Food Microbiol* 2008; **121(1)**: 41-52.

Habib I, Sampers I, Uyttendaele M, Berkvens D, De Zutter L. Performance characteristics and estimation of measurement uncertainty of three plating procedures for *Campylobacter* enumeration in chicken meat. *Food Microbiol* 2008; **25(1)**: 65-74.

Seal BS, Hiatt KL, Kuntz RL, Woolsey R, Schegg KM, Ard M, Stintzi A. Proteomic analyses of a robust versus a poor chicken gastrointestinal colonizing isolate of *Campylobacter jejuni*. *J Proteome Res* 2007; **6(12)**: 4582-91.

### ***Clostridium botulinum***

Tseng-Ong L, Mitchell WG. Infant botulism: 20 years' experience at a single institution. *J Child Neurol* 2007; **22(12)**: 1333-7.

Volland H, Lamourette P, Nevers MC, Mazuet C, Ezan E, Neuburger LM, Popoff M, Créminon C. A sensitive sandwich enzyme immunoassay for free or complexed *Clostridium botulinum* neurotoxin type A. J Immunol Methods 2008; **330(1-2)**: 120-9.

Gao YL, Ju XR. Exploiting the combined effects of high pressure and moderate heat with nisin on inactivation of *Clostridium botulinum* spores. J Microbiol Method 2008; **72(1)**: 20-8.

Bianco MI, Lúquez C, de Jong LI, Fernández RA. Presence of *Clostridium botulinum* spores in Matricaria chamomilla (chamomile) and its relationship with infant botulism. Int J Food Microbiol 2008; **121(3)**: 357-60.

### ***Clostridium perfringens***

Paredes- Sabja D, Torres JA, Setlow P, Sarker MR. *Clostridium perfringens* spore germination: characterization of germinants and their receptors. J Bacteriol 2008; **190(4)**: 1190-201.

Juneja VK, Marks H, Thippareddi H. Predictive model for growth of *Clostridium perfringens* during cooling of cooked uncured beef. Food Microbiol 2008; **25(1)**: 42-55.

Chalmers G, Martin SW, Hunter DB, Presscott JF, Weber LJ, Boerlin P. Genetic diversity *Clostridium perfringens* isolated from healthy broiler chickens at a commercial farm. Vet Microbiol 2008; **127(1-2)**: 116-27.

### ***Cryptosporidium***

Xiao L, Feng Y. Zoonotic cyptosporidiosis. FEMS Immunol Med Microbiol 2008 Jan 18 [Epub ahead of print].

Graczyk TK, Majewska AC, Schwab KJ. The role of birds in dissemination of human waterborne enteropathogens. Trends Parasitol 2008; **24(2)**: 55-59.

Starkey SR, White ME, Mohammed HO. Cryptosporidium and dairy cattle in the Catskill/Delaware watershed: a quantitative risk assessment. Risk Anal 2007; **27(6)**: 1469-85.

Brook EJ, Anthony Hart C, French NP, Christley RM. Molecular epidemiology of *Cryptosporidium* subtypes in cattle in England. Vet J 2007 Dec 11 [Epub ahead of print].

Leoni F, Gómez-Couso H, Ares-Mazás ME, McLauchlin J. Multilocus genetic analysis of *Cryptosporidium* in naturally contaminated bivalve molluscs. J Appl Microbiol 2007; **103(6)**: 2430-7.

Reynolds KA, Mena KD, Gerba CP. Risk of waterborne illness via drinking water in the United States. Rev Environ Contamin Toxicol 2008; **192**: 117-58. Review.

Kay D, Watkins J, Francis CA, Wyn-Jones AP, Stapleton CM, Fewtrell L, Wyer MD, Drury D. The microbiological quality of seven large commercial private water supplies in the United Kingdom. *J Water Health* 2007; **5(4)**: 523-38.

### ***E. Coli***

Jay MT, Cooley M, Carychao D, Wiscomb GW, Sweitzer RA, Crawford-Miksza L, Farrar JA, Lau DK, O'Connell J, Millington A, Asmundson RV, Atwill ER, Mandrell RE. *Escherichia coli* O157:H7 in feral swine near spinach fields and cattle, central California coast. *Emerg Infect Dis* 2007; **13(12)**: 1908-11.

Pradel N, Bertin Y, Martin C, Livrelli V. Molecular Analysis of Shiga Toxin-Producing *Escherichia coli* strains isolated from Hemolytic-Uremic Syndrome patients and dairy samples in France. *Appl Environ Microbiol* 2008 Feb 1 [Epub ahead of print].

Grant MA. Comparison of *Escherichia coli* O157:H7 enrichment in spiked produce samples. *J Food Prot* 2008. **71(1)**: 139-45.

Palanichamy A, Jayas DS, Holley RA. Predicting survival of *Escherichia coli* O157:H7 in dry fermented sausage using artificial neural networks. *J Food Prot* 2008;**71(1)**: 6(12).

Lavigne JP, Blanc-Potard AB. Molecular evolution of *Salmonella enterica* serovar Typhimurium and pathogenic *Escherichia coli*: From pathogenesis to therapeutics. *Infect Genet Evol* 2007 Nov 29 [Epub ahead of print]

Roest HI, Liebana E, Wannet W, Van Duynoven Y, Veldman KT, Mevius DJ. [Antibiotic resistance in *Escherichia coli* O157 isolated between 1998 and 2003 in The Netherlands.] *Tijdschr Diergeneeskd* 2007; **132(24)**: 954-8. Dutch.

Murphy M, Minihan D, Buckley JF, O'Mahony M, Whyte P, Fanning S. Multiple-locus variable number of tandem repeat analysis (MLVA) of Irish verocytotoxigenic *Escherichia coli* O157 from feedlot cattle: uncovering strain dissemination routes. *BMC Vet Res* 2008; **4**: 2

Semenov AV, Franz E, van Overbeek L, Termorshuizen AJ, van Bruggen AH. Estimating the stability of *Escherichia coli* O157:H7 survival in manure-amended soils with different management histories. *Environ Microbiol* 2008 Jan 21 [Epub ahead of print]

Joishy M, Jauhari P, Mathew AA, Rangarajan T. Can we predict the development of haemolytic uraemic syndrome in the early stage of *Escherichia coli* O157 infection? *Arch Dis Child* 2008; **93(2)**: 180-1.

Franz E, Semenov AV, Termorshuizen AJ, de Vos OJ, Bokhorst JG, van Bruggen AH. Manure-amended soil characteristics affecting the survival of *E. coli* O157:H7 in 36 Dutch soils. *Environ Microbiol* 2008; **10(2)**: 313-27.

Arnold ME, Ellis-Iversen J, Cook AJ, Davies RH, McLaren IM, Kay AC, Pritchard GC. Investigation into the effectiveness of pooled fecal samples for detection of verocytotoxin-producing *Escherichia coli* O157 in cattle. *J Vet Diagn Invest* 2008; **20(1)**: 21-7.

Iijima K, Kamioka I, Nozu K. Management of diarrhea-associated haemolytic uremic syndrome in children. *Clin Exp Nephrol* 2008; **12(1)**: 16-19.

Mukherjee A, Mammel MK, Leclerc JE, Cebula TA. Altered Utilization of N-Acetyl-D-Galactosamine by *Escherichia coli* O157:H7 from the 2006 Spinach Outbreak. *J Bacteriol* 2008; **190(5)**: 1710-7.

Martínez AJ, Bossio CP, Durango AC, Vanegas MC. Characterization of Shiga toxin-producing *Escherichia coli* isolated from foods. *J Food Prot* 2007; **70(12)**: 2843-6.

Shepherd MW Jr, Liang P, Jiang X, Doyle MP, Erickson MC. Fate of *Escherichia coli* O157:H7 during an on-farm dairy manure-based composting. *J Food Prot* 2007; **70(12)**: 2708-16.

Uhlich GA, Sinclair JR, Warren NG, Chmielecki WA, Fratamico P. Characterization of Shiga toxin-producing *Escherichia coli* isolates associated with two multistate food-borne outbreaks that occurred in 2006. *Appl Environ Microbiol* 2008; **74(4)**: 1268-72.

Ellis-Iversen J, Smith RP, Van Winden S, Paiba GA, Watson E, Snow LC, Cook AJ. Farm practices to control *E. coli* O157 in young cattle – a randomised controlled trial. *Vet Res* 2008; **39(1)**: 3.

Wu G, Carter B, Mafura M, Liebana E, Woodward MJ, Anjum MF. Genetic diversity among *Escherichia coli* O157:H7 isolates and identification of genes linked to human infections. *Infect Immun* 2008; **76(2)**: 845-56.

Bettelheim K. Re: Isolation of *Escherichia coli* O157:H7 strains that do not produce Shiga toxin from bovine, avian and environmental sources. *Lett Appl Microbiol* 2008; **46(2)**: 281.

Solecki O, MacRae M, Ogden I, Strachan N. Can the high levels of human verocytotoxigenic *Escherichia coli* O157 infection in rural areas of NE Scotland be explained by consumption of contaminated meat? *J Appl Microbiol* 2007; **103(6)**: 2616-21.

Smet A, Martel A, Persoons D, Dewulf J, Heyndrickx M, Catry B, Herman L, Haesebrouck F, Butaye P. Diversity of extended-spectrum-beta-lactamases and class C beta-lactamases among cloacal *Escherichia coli* in Belgian broiler farms. *Antimicrob Agents Chemother* 2008 Jan 28; [Epub ahead of print]

Meyer E, Lunke C, Kist M, Schwab F, Frank U. Antimicrobial Resistance in *Escherichia coli* Strains Isolated from Food, Animals and Humans in Germany. *Infection* 2008; **36(1)**: 59-61.

Nicolas-Chanoine MH, Blanco J, Leflon-Guibout V, Demarty R, Alonso MP, Caniça MM, Park YJ, Lavigne JP, Pitout J, Johnson JR. Intercontinental emergence of *Escherichia coli* clone O25:H4-ST131 producing CTX-M-15. *J Antimicrob Chemother* 2008; **61(2)**: 273-81.

Karisik E, Ellington MJ, Livermore DM, Woodford N. Virulence factors in *Escherichia coli* with CTX-M-15 and other extended-spectrum beta-lactamases in the UK. *J Antimicrob Chemother* 2008; **61(1)**: 54-8.

### ***Hepatitis A***

Papafragkou E, Plante M, Mattison K, Bidawid S, Karthikeyan K, Farber JM, Jaykus LA. Rapid and sensitive detection of hepatitis A virus in representative food matrices. *J Virol Methods* 2008; **147(1)**: 177-87.

Casas N, Amarita F, de Marañón IM. Evaluation of an extracting method for the detection of Hepatitis A virus in shellfish by SYBR-Green real-time RT-PCR. *Int J Food Microbiol* 2007; **120(1-2)**: 179-85.

### ***Hepatitis E***

Yu JN, Kim MY, Kim DG, Kim SE, Lee JB, Park SY, Song CS, Shin HC, Seo KH, Choi IS. Prevalence of hepatitis E virus and sapovirus in post-weaning pigs and identification of their genetic diversity. *Arch Virol* 2008 Feb 9 [Epub ahead of print]

Lewis HC, Boisson S, Ijaz S, Hewitt K, Ngui SL, Boxall E, Teo CG, Morgan D. Hepatitis E in England and Wales. *Emerg Infect Dis* 2008; **14(1)**: 165-7.

Purcell RH, Emerson SU. Hepatitis E: An emerging awareness of an old disease. *J Hepatol* 2008; **48(3)**: 494-503.

Deest G, Zehner L, Nicand E, Gaudy-Graffin C, Goudeau A, Bacq Y. [Autochthonous hepatitis E in France and consumption of raw pig meat.] *Gastroenterol Clin Biol* 2007; **31(12)**: 1095-7. French.

Feagins AR, Opriessnig T, Guenette DK, Halbur PG, Meng XJ. Inactivation of infectious hepatitis E virus present in commercial pig livers sold in local grocery stores in the United States. *Int J Food Microbiol* 2007; Dec 4 [Epub ahead of print]

De Silva AN, Muddu AK, Iredale JP, Sheron N, Khakoo SI, Pelosi E. Unexpectedly high incidence of indigenous acute hepatitis E within South Hampshire: time for routine testing? *J Med Virol* 2008; **80(2)**: 283-8.



Bouwknegt M, Lodder-Verschoor F, van der Poel WH, Rutjes SA, de Roda Husman AM. Hepatitis E virus RNA in commercial porcine livers in The Netherlands. *J Food Prot* 2007; **70(12)**: 2889-95.

### ***Listeria monocytogenes***

Porsby CH, Vogel VG, Mohr M, Gram L. Influence of processing steps in cold-smoked salmon production on survival and growth of persistent and presumed non-persistent *Listeria monocytogenes*. *Int J Food Microbiol* 2008 Jan 26 [Epub ahead of print].

Guilbaud M, Chafsey I, Pilet MF, Leroi F, Prévost H, Hébraud M, Dousset X. Response of *Listeria monocytogenes* to liquid smoke. *J Appl Microbiol* 2008 Feb 6; [Epub ahead of print].

Adrião A, Vieira M, Fernandes I, Barbosa M, Sol M, Tenreiro RP, Chambel L, Barata B, Zilhao I, Sharma G, Perni S, Jordan SJ, Andrew PW, Faleiro ML. Marked intra-strain variation in response of *Listeria monocytogenes* dairy isolates to acid or salt stress and the effect of acid or salt adaptation on adherence to abiotic surfaces. *Int J Food Microbiol* 2007 Dec 28 [Epub ahead of print].

Sperry KE, Kathariou S, Edwards JS, Wolf LA. Multiple-Locus Variable Number Tandem Repeat Analysis as a subtyping tool for *Listeria monocytogenes*. *J Clin Microbiol* 2008 Feb 6 [Epub ahead of print]

Abadias M, Usall J, Oliveira M, Alegre I, Viñas I. Efficacy of neutral electrolyzed water (NEW) for reducing microbial contamination on minimally-processed vegetables. *Int J Food Microbiol* 2008; Jan 29 [Epub ahead of print].

Keskinen LA, Todd EC, Ryser ET. Transfer of surface-dried *Listeria monocytogenes* from stainless steel knife blades to roast turkey breast. *J Food Prot* 2008; **71(1)**: 176-81.

Gougouli M, Angelidis AS, Koutsoumanis K. A study on the kinetic behaviour of *Listeria monocytogenes* in ice cream stored under static and dynamic chilling and freezing conditions. *J Dairy Sci* 2008; **91(2)**: 523-30.

Mullapudi S, Siletzky RM, Kathariou S. Heavy metal and bezalkonium chloride resistance of *Listeria monocytogenes* from the environment of turkey processing plants. *Appl Environ Microbiol* 2008; Jan 11 [Epub ahead of print].

Lundén J, Tolvanen R, Korkeala H. Acid and heat tolerance of persistent and nonpersistent *Listeria monocytogenes* food plant strains. *Lett Appl Microbiol* 2008; **46(2)**: 276-80.

Navas J, Ortiz S, López V, Martínez-Suárez JV. Different enrichment procedures for recovery of *Listeria monocytogenes* from raw chicken samples

can affect the results of detection (by chromogenic plating or real-time PCR) and lineage or strain identification. J Food Prot 2007; **70(12)**: 2851-4.

Smith MA, Takeuchi K, Anderson G, Ware GO, McClure HM, Raybourne RB, Mytle N, Doyle MP. Dose-response model for *Listeria monocytogenes*-induced stillbirths in nonhuman primates. Infect Immun 2008; **76(2)**: 726-31.

Rantsiou K, Alessandria V, Urso R, Dolci P, Cocolin L. Detection, quantification and vitality of *Listeria monocytogenes* in food as determined by quantitative PCR. Int J Food Microbiol 2008; **121(1)**: 99-105

Guentzel JL, Liang Lam K, Callan MA, Emmons SA, Dunham VL. Reduction of bacteria on spinach, lettuce, and surfaces in food service areas using neutral electrolyzed oxidising water. Food Microbiol 2008; **25(1)**: 36-41.

Oravcová K, Trncíková T, Kuchta T, Kaclíková E. Limitation in the detection of *Listeria monocytogenes* in food in the presence of competing *Listeria innocua*. J Appl Microbiol 2008; **104(2)**: 429-37.

De Cesare A, Mioni R, Manfreda G. Prevalence of *Listeria monocytogenes* in fresh and fermented Italian sausages and ribotyping of contaminating strains. Int J Food Microbiol 2007; **120(1-2)**: 124-30.

### **Norovirus**

Lopman B, Zambon M, Brown DW. The Evolution of Norovirus, the “Gastric Flu” PLoS Med 2008; **5(2)**: e42. [Epub ahead of print].

Lindesmith LC, Donaldson EF, Lobue AD, Cannon JL, Zheng DP, Vinje J, Baric RS. Mechanisms of GII.4 Norovirus Persistence in Human Populations. PLoS Med 2008; **5(2)**: e31. [Epub ahead of print].

Baert L, Uyttendaele M, Debevere J. Evaluation of viral extraction methods on a broad range of Ready-to-Eat foods with conventional and real-time RT-PCR for Norovirus GII detection. Int J Food Microbiol 2008; Jan 2 [Epub ahead of print]

Kelly S, Foley B, Dunford L, Coughlan S, Tuite G, Duffy M, Mitchell S, Smyth B, O'Neill H, McKeown P, Hall W, Lynch M. Establishment of a national database to link epidemiological and molecular data from norovirus outbreaks in Ireland. Epidemiol Infect 2008; Feb 6; 1-8 [Epub ahead of print].

Lee J, Zoh K, Ko G. Inactivation and UV/TiO<sub>2</sub> Disinfection of Murine Norovirus under Various Environmental Conditions. Appl Environ Microbiol 2008 Feb 1 [Epub ahead of print].

Noda M, Fukuda S, Nishio O. Statistical analysis of attack rate in norovirus foodborne outbreaks. Int J Food Microbiol 2008; **122(1-2)**: 216-20.

Showell D, Sundkvist T, Reacher M, Gray J. Norovirus outbreak associated with canteen salad in Suffolk, United Kingdom. *Euro Surveill* 2007; **12(11)**: E071129.6.

Ozawa K, Oka T, Takeda N, Hansman GS. Norovirus infections in symptomatic and asymptomatic food handlers in Japan. *J Clin Microbiol* 2007; **45(12)**: 3996-4005.

Iturriza Gómara M, Simpson R, Perault AM, Redpath C, Lorgelly P, Joshi D, Mugford M, Hughes CA, Dalrymple J, Desselberger U, Gray J. Structured surveillance of infantile gastroenteritis in East Anglia, UK: incidence of infection with common viral gastroenteric pathogens. *Epidemiol Infect* 2008; **136(1)**: 23-33.

### ***Salmonella***

Perron GG, Bell G, Quessy S. Parallel evolution of multidrug-resistance in *Salmonella enterica* isolated from swine. *FEMS Microbiol Lett* 2008 Feb 12 [Epub ahead of print]

Townes JM, Deodhar AA, Laine ES, Smith K, Krug HE, Barkhuizen A, Thompson ME, Cieslak PR, Sobel J. Reactive arthritis following culture-confirmed infections with bacterial enteric pathogens in Minnesota and Oregon: a population-based study. *Ann Rheum Dis* 2008 Feb 13 [Epub ahead of print]

OzFoodNet Working Group. Monitoring the incidence and causes of diseases potentially transmitted by food in Australia: annual report of the Ozfoodnet Network, 2006. *Commun Dis Intell* 2007; **31(4)**: 345-65.

Prévost K, Magal P, Protais J, Beaumont C. Effect of genetic resistance of the hen to *Salmonella* carrier-state on incidence of bacterial contamination: synergy with vaccination. *Vet Res* 2008; **39(2)**:20.

Hopkins KL, Day M, Threlfall EJ. Plasmid-mediated Quinolone Resistance in *Salmonella enterica*, United Kingdom. *Emerg Infect Dis* 2008; **14(2)**: 340-2.

Hedberg CW, Greenblatt JF, Matyas BT, Lemmings J, Sharp DJ, Skibicki RT, Liang AP, Timeline Study Work Group TE. Timeliness of Enteric Disease Surveillance in 6 US States. *Emerg Infect Dis* 2008; **14(2)**: 311-3.

Ternhag A, Törner A, Svensson A, Ekdahl K, Giesecke J. Short- and long-term effects of bacterial gastrointestinal infections. *Emerg Infect Dis* 2008; **14(1)**: 143-8.

Zheng J, Keys CE, Zhao S, Meng J, Brown EW. Enhanced subtyping scheme for *Salmonella enteritidis*. *Emerg Infect Dis* 2007; **13(12)**: 1932-5.

Heithoff DM, Shimp WR, Lau PW, Badie G, Enioutina EY, Daynes RA, Byrne BA, House JK, Mahan MJ. Human *Salmonella* Clinical Isolates That Are

Distinct From Those of Animal Origin. *Appl Environ Microbiol* 2008 Feb 1 [Epub ahead of print].

Boyen F, Haesebrouck F, Maes D, Van Immerseel F, Ducatelle R, Pasmans F. Non-typhoidal *Salmonella* infections in pigs: A closer look at epidemiology, pathogenesis and control. *Vet Microbiol* 2008 Jan 4 [Epub ahead of print].

Abadias M, Usall J, Anguera M, Solsona C, Viñas I. Microbiological quality of fresh, minimally-processed fruit and vegetables, and sprouts from retail establishments. *Int J Food Microbiol* 2008 Jan 29 [Epub ahead of print]

Little CL, Rhoades JR, Hucklesby L, Greenwood M, Surman-Lee S, Bolton FJ, Meldrum R, Wilson I, McDonald C, de Pinna E, Threlfall EJ, Chan CH. Survey of *Salmonella* contamination of raw shell eggs used in food service premises in the United Kingdom, 2005 through 2006. *J Food Prot* 2008; **71(1)**: 19-26.

Esteban JI, Oporto B, Aduriz G, Juste RA, Hurtado A. A survey of food-borne pathogens in free-range poultry farms. *Int J Food Microbiol* 2008; Jan 28 [Epub ahead of print].

Philbey AW, Mather HA, Taylor DJ, Coia JE. Isolation of avian strains of *Salmonella enterica* serovar Typhimurium from cats with enteric disease in the United Kingdom. *Vet Rec* 2008; **162(4)**: 120-2.

Lang NL, Smith SR. Time and temperature inactivation kinetics of enteric bacteria relevant to sewage sludge treatment processes for agricultural use. *Water Res* 2007; Dec 14 [Epub ahead of print].

Benschop J, Stevenson MA, Dahl J, Morris RS, French NP. Temporal and longitudinal analysis of Danish Swine Salmonellosis Control Programme data: implications for surveillance. *Epidemiol Infect* 2008 Jan 16;: 1-10 [Epub ahead of print].

Marcus R. New information about pediatric foodborne infections: the view from FoodNet. *Curr Opin Pediatr* 2008; **20(1)**: 79-84.

Denny J, Threlfall J, Takkinen J, Lofdahl S, Westrell T, Varela C, Adak B, Boxall N, Ethelberg S, Torpdahl M, Straetemans M, van Pelt W. Multinational *Salmonella* Paratyphi B variant Java (*Salmonella* Java) outbreak, August – December 2007. *Euro Surveill* 2007; **12(12)**: E071220.2.

Jenkins MB, Endale DM, Fisher DS. Most probable number methodology for quantifying dilute concentrations and fluxes of *Salmonella* in surface waters. *J Appl Microbiol* 2007 Dec 20 [Epub ahead of print].

Yoke-Kqueen C, Learn-Han L, Noorzaleha AS, Son R, Sabrina S, Jiun-Horng S, Chai-Hoon K. Characterization of multiple-antimicrobial-resistant *Salmonella enterica* Subsp. *enterica* isolated from indigenous vegetables and poultry in Malaysia. *Lett Appl Microbiol* 2007 Dec 20 [Epub ahead of print].

## ***Toxoplasma***

Yolken RH, Torrey EF. Are some cases of psychosis caused by microbial agents? A review of the evidence. *Mol Psychiatry* 2008 Feb 12 [Epub ahead of print].

Schares G, Vrhovec MG, Pantchev N, Herrmann DC, Conraths FJ. Occurrence of *Toxoplasma gondii* and *Hammondia hammondi* oocysts in the faeces of cats from Germany and other European countries. *Vet Parasitol* 2008; **152(1-2)**: 34-45.

Dubey JP, Sundar N, Hill D, Velmurugan GV, Bandini LA, Kwok OC, Majumdar D, Su C. High prevalence and abundant atypical genotypes of *Toxoplasma gondii* isolated from lambs destined for human consumption in the USA. *Int J Parasitol* 2007 Dec 8 [Epub ahead of print].

Moré G, Basso W, Bacigalupe D, Venturini MC, Venturini L. Diagnosis of *Sarcocystis cruzi*, *Neospora caninum*, and *Toxoplasma gondii* infections in cattle. *Parasitol Res* 2008; **102(4)**: 671-5.

Dabritz HA, Miller MA, Atwill ER, Gardner IA, Leutenegger CM, Melli AC, Conrad PA. Detection of *Toxoplasma gondii*-like oocysts in cat feces and estimates of the environmental oocyst burden. *J Am Vet Med Assoc* 2007; **231(11)**: 1676-84.

Correa D, Cañedo-Solares I, Ortiz-Alegría LB, Caballero-Ortega H, Rico-Torres CP. Congenital and acquired toxoplasmosis: diversity and role of antibodies in different compartments of the host. *Parasite Immunol* 2007; **29(12)**: 651-60. Review.

Gollub EL, Leroy V, Gilbert R, Chêne G, Wallon M; the European Toxoprevention Study Group (EUROTOXO). Effectiveness of health education on *Toxoplasma*-related knowledge, behaviour, and risk of seroconversion in pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2008; **136(2)**: 137-145.

Robertson LJ. The potential for marine bivalve shellfish to act as transmission vehicles for outbreaks of protozoan infections in humans: a review. *Int J Food Microbiol* 2007; **120(3)**: 201-16.