

ADVISORY COMMITTEE ON THE MICROBIOLOGICAL SAFETY OF FOOD

INFORMATION PAPER

Items of interest from the literature***Bacillus cereus***

Daelman J, Membré JM, Jacxsens L, Vermeulen A, Devlieghere F, Uyttendaele M. A quantitative microbiological exposure assessment model for *Bacillus cereus* in REPFEDs. *Int J Food Microbiol.* 2013;166(3): 433-449.

Eijlander RT, Kuipers OP. Live-Cell Imaging Tool Optimization To Study Gene Expression Levels and Dynamics in Single Cells of *Bacillus cereus*. *Appl Environ Microbiol.* 2013; 79(18):5643-51.

Daelman J, Sharma A, Vermeulen A, Uyttendaele M, Devlieghere F, Membré JM. Development of a time-to-detect growth model for heat-treated *Bacillus cereus* spores. *Int J Food Microbiol.* 2013;165(3): 231-40.

Ceuppens S, Boon N, Uyttendaele M. Diversity of *Bacillus cereus* group strains is reflected in their broad range of pathogenicity and diverse ecological lifestyles. *FEMS Microbiol Ecol.* 2013; 84(3): 433-50.

Campylobacter

Aroori SV, Cogan TA, Humphrey TJ. The effect of growth temperature on the pathogenicity of campylobacter. *Curr Microbiol.* 2013; 67(3): 333-40.

Bouwknegt M, van Pelt W, Havelaar AH. Scoping the impact of changes in population age-structure on the future burden of foodborne disease in the Netherlands, 2020-2060. *Int J Environ Res Public Health.* 2013; 10(7): 2888-96.

Chen Y, Mukherjee S, Hoffmann M, Kotewicz ML, Young S, Abbott J, Luo Y, Davidson MK, Allard M, McDermott P, Zhao S. Whole Genome Sequencing of a Gentamicin-Resistant *Campylobacter coli* isolated from United States Retail Meats Reveals Novel Plasmid Mediated Aminoglycoside Resistance Genes. *Antimicrob Agents Chemother.* 2013 Aug 19. [Epub ahead of print] PubMed PMID: 23959310.

Chon JW, Kim H, Kim HS, Seo KH. Improvement of modified charcoal-cefoperazone-deoxycholate agar by addition of potassium clavulanate for detecting *Campylobacter* spp. in chicken carcass rinse. *Int J Food Microbiol.* 2013; 165(1): 7-10.

Ge B, Wang F, Sjölund-Karlsson M, McDermott PF. Antimicrobial resistance in *Campylobacter*: Susceptibility testing methods and resistance trends. *J Microbiol Methods*. 2013 ; 95(1): 57-67.

Gharst G, Oyarzabal OA, Hussain SK. Review of current methodologies to isolate and identify *Campylobacter* spp. from foods. *J Microbiol Methods*. 2013; 95(1): 84-92.

Gilpin BJ, Walsh G, On SL, Smith D, Marshall JC, French NP. Application of molecular epidemiology to understanding campylobacteriosis in the Canterbury region of New Zealand. *Epidemiol Infect*. 2013;141(6): 1253-66.

Guyard-Nicodème M, Tresse O, Houard E, Jugiau F, Courtillon C, El Manaa K, Laisney MJ, Chemaly M. Characterization of *Campylobacter* spp. transferred from naturally contaminated chicken legs to cooked chicken slices via a cutting board. *Int J Food Microbiol*. 2013; 164(1): 7-14.

Hardy B, Crilly N, Pendleton S, Andino A, Wallis A, Zhang N, Hanning I. Impact of rearing conditions on the microbiological quality of raw retail poultry meat. *J Food Sci*. 2013; 78(8): M1232-5.

Hu Y, Shang Y, Huang J, Wang Y, Ren F, Jiao Y, Pan Z, Jiao XA. A novel immunoproteomics method for identifying in vivo-induced *Campylobacter jejuni* antigens using pre-adsorbed sera from infected patients. *Biochim Biophys Acta*. 2013; 1830(11): 5229-5235.

Isohanni P, Huehn S, Aho T, Alter T, Lyhs U. Heat stress adaptation induces cross-protection against lethal acid stress conditions in *Arcobacter butzleri* but not in *Campylobacter jejuni*. *Food Microbiol*. 2013; 34(2): 431-5.

Klančnik A, Vučković D, Plankl M, Abram M, Smole Možina S. *In vivo* modulation of *Campylobacter jejuni* virulence in response to environmental stress. *Foodborne Pathog Dis*. 2013; 10(6): 566-72.

Kudirkiene E, Bunevičienė J, Šernienė L, Ramonaitė S, Olsen JE, Malakauskas M. Importance of the producer on retail broiler meat product contamination with *Campylobacter* spp. *J Sci Food Agric*. 2013; 93(9): 2293-8.

Lake RJ, Horn BJ, Dunn AH, Parris R, Green FT, McNickle DC. Cost-effectiveness of interventions to control *Campylobacter* in the New Zealand poultry meat food supply. *J Food Prot*. 2013; 76(7): 1161-7.

Meredith H, Walsh D, McDowell DA, Bolton DJ. An investigation of the immediate and storage effects of chemical treatments on *Campylobacter* and sensory characteristics of poultry meat. *Int J Food Microbiol.* 2013; 166(2): 309-15.

Mughini-Gras L, Smid JH, Wagenaar JA, DE Boer A, Havelaar AH, Friesema IH, French NP, Graziani C, Busani L, Van Pelt W. *Campylobacteriosis* in returning travellers and potential secondary transmission of exotic strains. *Epidemiol Infect.* 2013: 1-12. [Epub ahead of print]

On SL. Isolation, identification and subtyping of *Campylobacter*: Where to from here? *J Microbiol Methods.* 2013; 95(1): 3-7.

Schijven J, Bouwknegt M, de Roda Husman AM, Rutjes S, Sudre B, Suk JE, Semenza JC. A Decision Support Tool to Compare Waterborne and Foodborne Infection and/or Illness Risks Associated with Climate Change. *Risk Anal.* 2013 Jun 19. doi:10.1111/risa.12077. [Epub ahead of print]

Schmidt PJ, Pintar KD, Fazil AM, Flemming CA, Lanthier M, Laprade N, Sunohara MD, Simhon A, Thomas JL, Topp E, Wilkes G, Lapen DR. Using *Campylobacter* spp. And *Escherichia coli* data and Bayesian microbial risk assessment to examine public health risks in agricultural watersheds under tile drainage management. *Water Res.* 2013; 47(10): 3255-72.

Ugarte-Ruiz M, Wassenaar TM, Gómez-Barrero S, Porrero MC, Navarro-Gonzalez N, Domínguez L. The effect of different isolation protocols on detection and molecular characterization of *Campylobacter* from poultry. *Lett Appl Microbiol.* 2013 Jul 10. doi: 10.1111/lam.12130. [Epub ahead of print]

Wei S, Gutek A, Lilburn M, Yu Z. Abundance of pathogens in the gut and litter of broiler chickens as affected by bacitracin and litter management. *Vet Microbiol.* 2013; 166(3-4): 595-601.

Wieczorek K, Kania I, Osek J. Prevalence and Antimicrobial Resistance of *Campylobacter* spp. Isolated from Poultry Carcasses in Poland. *J Food Prot.* 2013; 76(8):1451-5.

Wieczorek K, Osek J. Antimicrobial resistance mechanisms among *Campylobacter*. *Biomed Res Int.* 2013; 2013:340605. doi: 10.1155/2013/340605. Epub 2013 Jun 24.

Clostridium

André S, Zuber F, Remize F. Thermophilic spore-forming bacteria isolated from spoiled canned food and their heat resistance. Results of a French ten-year survey. *Int J Food Microbiol.* 2013; 165(2):134-43.

Bennett SD, Walsh KA, Gould LH. Foodborne disease outbreaks caused by *Bacillus cereus*, *Clostridium perfringens*, and *Staphylococcus aureus*--United States, 1998-2008. *Clin Infect Dis.* 2013; 57(3): 425-33.

Bjornsdottir-Butler K, McCarthy S, Burkhardt W 3rd, Benner RA Jr. Importance of histamine-producing *Clostridium perfringens* in scombrotoxin-forming fish. *J Food Prot.* 2013; 76(7): 1283-7.

Derman Y, Isokallio M, Lindström M, Korkeala H. The two-component system CBO2306/CBO2307 is important for cold adaptation of *Clostridium botulinum* ATCC 3502. *Int J Food Microbiol.* 2013 Jun 14. doi:pii: S0168-1605(13)00291-2. 10.1016/j.ijfoodmicro.2013.06.004. [Epub ahead of print]

Eckert C, Burghoffer B, Barbut F. Contamination of ready-to-eat raw vegetables with *Clostridium difficile* in France. *J Med Microbiol.* 2013; 62(Pt 9): 1435-8.

Guran HS, Oksuztepe G. Detection and typing of *Clostridium perfringens* from retail chicken meat parts. *Lett Appl Microbiol.* 2013; 57(1): 77-82.

Hoover DG, Rodriguez-Palacios A. Transmission of *Clostridium difficile* in Foods. *Infect Dis Clin North Am.* 2013; 27(3): 675-85.

Li T, Tian R, Cai K, Wang Q, Chen F, Fang H, Luo S, Li Z, Wang D, Hou X, Wang H. The Effect of pH on Growth of *Clostridium botulinum* Type A and Expression of bontA and botR During Different Growth Stages. *Foodborne Pathog Dis.* 2013; 10(8): 692-7.

Malakar PK, Plowman J, Aldus CF, Xing Z, Zhao Y, Peck MW. Detection limit of *Clostridium botulinum* spores in dried mushroom samples sourced from China. *Int J Food Microbiol.* 2013; 166(1): 72-6.

Marsh JW. Counterpoint: Is *Clostridium difficile* a food-borne disease? *Anaerobe.* 2013; 21: 62-3.

Ohtani K, Hirakawa H, Paredes-Sabja D, Tashiro K, Kuhara S, Sarker MR, Shimizu T. Unique regulatory mechanism of sporulation and enterotoxin production in *Clostridium perfringens*. *J Bacteriol.* 2013; 195(12): 2931-6.

Reddy NR, Marshall KM, Morrissey TR, Loeza V, Patazca E, Skinner GE, Krishnamurthy K, Larkin JW. Combined High Pressure and Thermal Processing on Inactivation of Type A and Proteolytic Type B Spores of *Clostridium botulinum*. J Food Prot. 2013; 76(8): 1384-92.

Redondo-Solano M, Valenzuela-Martinez C, Cassada DA, Snow DD, Juneja VK, Burson DE, Thippareddi H. Effect of meat ingredients (sodium nitrite and erythorbate) and processing (vacuum storage and packaging atmosphere) on germination and outgrowth of *Clostridium perfringens* spores in ham during abusive cooling. Food Microbiol. 2013; 35(2): 108-15.

Rodriguez-Palacios A, Borgmann S, Kline TR, LeJeune JT. *Clostridium difficile* in foods and animals: history and measures to reduce exposure. Anim Health Res Rev. 2013; 14(1): 11-29.

Udompijikul P, Alnoman M, Sarker MR. Inactivation strategy for *Clostridium perfringens* spores adhered to food contact surfaces. Food Microbiol. 2013; 34(2): 328-36.

Wei S, Gutek A, Lilburn M, Yu Z. Abundance of pathogens in the gut and litter of broiler chickens as affected by bacitracin and litter management. Vet Microbiol. 2013; 166(3-4): 595-601.

Cryptosporidium

Hohweyer J, Dumètre A, Aubert D, Azas N, Villena I. Tools and methods for detecting and characterizing giardia, cryptosporidium, and toxoplasma parasites in marine mollusks. J Food Prot. 2013; 76(9): 1649-57.

Jex AR, Koehler AV, Ansell BR, Baker L, Karunajeewa H, Gasser RB. Getting to the guts of the matter: The status and potential of 'omics' research of parasitic protists of the human gastrointestinal system. Int J Parasitol. 2013 Jul 18. doi:pii: S0020-7519(13)00177-X. 10.1016/j.ijpara.2013.06.005. [Epub ahead of print]

Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, Wu Y, Sow SO, Sur D, Breiman RF, Faruque AS, Zaidi AK, Saha D, Alonso PL, Tamboura B, Sanogo D, Onwuchekwa U, Manna B, Ramamurthy T, Kanungo S, Ochieng JB, Omere R, Oundo JO, Hossain A, Das SK, Ahmed S, Qureshi S, Quadri F, Adegbola RA, Antonio M, Hossain MJ, Akinsola A, Mandomando I, Nhampossa T, Acácio S, Biswas K, O'Reilly CE, Mintz ED, Berkeley LY, Muhsen K, Sommerfelt H, Robins-Browne RM, Levine MM. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. Lancet. 2013; 382(9888): 209-22.

Manser M, Granlund M, Edwards H, Saez A, Petersen E, Evengard B, Chiodini P; The European Society of Clinical Microbiology and Infectious Diseases Study Group on Clinical Parasitology. Detection of Cryptosporidium and Giardia in clinical laboratories in Europe-a comparative study. Clin Microbiol Infect. 2013 Jun 18. doi: 10.1111/1469-0691.12297. [Epub ahead of print] PubMed PMID: 24033667.

Marcos LA, Gotuzzo E. Intestinal protozoan infections in the immunocompromised host. Curr Opin Infect Dis. 2013; 26(4): 295-301.

McBride GB, Stott R, Miller W, Bambic D, Wuertz S. Discharge-based QMRA for estimation of public health risks from exposure to stormwater-borne pathogens in recreational waters in the United States. Water Res. 2013; 47(14): 5282-97.

Slapeta J. Cryptosporidiosis and Cryptosporidium species in animals and humans: A thirty colour rainbow? Int J Parasitol. 2013 Aug 20. doi:pii: S0020-7519(13)00202-6. 10.1016/j.ijpara.2013.07.005. [Epub ahead of print]

E.coli

Alhelfi NA, Adam H, Jones DL, Williams AP. Absence of *E coli* O157:H7 in sheep and cattle faeces in North Wales. Vet Rec. 2013; 173(6): 143.

Bagordo F, Grassi T, Idolo A, Serio F, Gabutti G, De Donno A. Rotavirus Occurrence in Shellfish with Low Levels of *E. coli*. Food Environ Virol. 2013 Jul 30. [Epub ahead of print] PubMed PMID: 23897404.

Beutin L, Hammerl JA, Reetz J, Strauch E. Shiga toxin-producing *Escherichia coli* strains from cattle as a source of the Stx2a bacteriophages present in enteroaggregative *Escherichia coli* O104:H4 strains. Int J Med Microbiol. 2013 Aug 16. doi:pii: S1438-4221(13)00107-0. 10.1016/j.ijmm.2013.08.001. [Epub ahead of print]

Bolton DJ, Ennis C, McDowell D. Occurrence, Virulence Genes and Antibiotic Resistance of Enteropathogenic *Escherichia coli* (EPEC) from Twelve Bovine Farms in the North-East of Ireland. Zoonoses Public Health. 2013 Jun 19. doi: 10.1111/zph.12058. [Epub ahead of print].

Call DR, Matthews L, Subbiah M, Liu J. Do antibiotic residues in soils play a role in amplification and transmission of antibiotic resistant bacteria in cattle populations? Front Microbiol. 2013; 4:193.

Comery R, Thanabalasuriar A, Garneau P, Portt A, Boerlin P, Reid-Smith RJ, Harel J, Manges AR, Gruenheid S. Identification of potentially diarrheagenic atypical

enteropathogenic *Escherichia coli* strains present in Canadian food animals at slaughter and in retail meats. *Appl Environ Microbiol.* 2013; 79(12): 3892-6.

Elkins CA, Kotewicz ML, Jackson SA, Lacher DW, Abu-Ali GS, Patel IR. Genomic paradigms for food-borne enteric pathogen analysis at the USFDA: case studies highlighting method utility, integration and resolution. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess.* 2013; 30(8): 1422-36.

Gayán E, Mañas P, Álvarez I, Condón S. Mechanism of the synergistic inactivation of *Escherichia coli* by UV-C light at mild temperatures. *Appl Environ Microbiol.* 2013; 79(14): 4465-73.

Gueimonde M, Sánchez B, G de Los Reyes-Gavilán C, Margolles A. Antibiotic resistance in probiotic bacteria. *Front Microbiol.* 2013;4:202.

Guernec A, Robichaud-Rincon P, Saucier L. Whole-genome transcriptional analysis of *Escherichia coli* during heat inactivation processes related to industrial cooking. *Appl Environ Microbiol.* 2013 Aug;79(16):4940-50.

Holvoet K, Sampers I, Callens B, Dewulf J, Uyttendaele M. Moderate prevalence of antimicrobial resistance in *E. coli* isolated from lettuce, irrigation water and soil. *Appl Environ Microbiol.* 2013 Aug 23. [Epub ahead of print] PubMed PMID: 23974140.

Hou Z, Fink RC, Sugawara M, Diez-Gonzalez F, Sadowsky MJ. Transcriptional and functional responses of *Escherichia coli* O157:H7 growing in the lettuce rhizosphere. *Food Microbiol.* 2013; 35(2): 136-42.

Händel N, Schuurmans JM, Brul S, ter Kuile BH. Compensation of the metabolic costs of antibiotic resistance by physiological adaptation in *Escherichia coli*. *Antimicrob Agents Chemother.* 2013; 57(8): 3752-62.

Klumpp J, Fouts DE, Sozhamannan S. Bacteriophage functional genomics and its role in bacterial pathogen detection. *Brief Funct Genomics.* 2013; 12(4): 354-65.

Kothari C, Gaiind R, Singh LC, Sinha A, Kumari V, Arya S, Chellani H, Saxena S, Deb M. Community acquisition of β -lactamase producing Enterobacteriaceae in neonatal gut. *BMC Microbiol.* 2013; 13: 136.

Lorenz SC, Son I, Maounounen-Laasri A, Lin A, Fischer M, Kase JA. Prevalence of Hemolysin Genes and Comparison of ehxA Subtype Patterns in Shiga toxin-producing *Escherichia coli* (STEC) and non-STEC Strains from Clinical, Food, and Animal Sources. *Appl Environ Microbiol.* 2013 Aug 9. [Epub ahead of print] PubMed PMID: 23934487.

Mellata M. Human and Avian Extraintestinal Pathogenic *Escherichia coli*: Infections, Zoonotic Risks, and Antibiotic Resistance Trends. *Foodborne Pathog Dis.* 2013 Aug 20. [Epub ahead of print] PubMed PMID: 23962019.

Määttä J, Lehto M, Kuisma R, Kymäläinen HR, Mäki M. Microbiological quality of fresh-cut carrots and process waters. *J Food Prot.* 2013; 76(7): 1240-4.

Page AV, Liles WC. Enterohemorrhagic *Escherichia coli* Infections and the Hemolytic-Uremic Syndrome. *Med Clin North Am.* 2013; 97(4): 681-95, xi.

Peng S, Hummerjohann J, Stephan R, Hammer P. Short communication: heat resistance of *Escherichia coli* strains in raw milk at different subpasteurization conditions. *J Dairy Sci.* 2013; 96(6): 3543-6.

Platteel TN, Leverstein-Van Hall MA, Cohen Stuart JW, Voets GM, van den Munckhof MP, Scharringa J, van de Sande N, Fluit AC, Bonten MJ; ESBL National Surveillance Working Group. Differences in the antibiotic susceptibility of human *Escherichia coli* with poultry-associated and non-poultry-associated extended-spectrum beta-lactamases. *Eur J Clin Microbiol Infect Dis.* 2013; 32(8): 1091-5.

Reeves HE, Lotz SB, Kennedy E, Randall LP, Coldham NG, La Ragione RM. Evaluation of an autogenous vaccine in cattle against *Escherichia coli* bearing the CTX-M-14 plasmid. *Res Vet Sci.* 2013; 94(3): 419-24.

Reich F, Atanassova V, Klein G. Extended-spectrum β -lactamase- and AmpC-producing enterobacteria in healthy broiler chickens, Germany. *Emerg Infect Dis.* 2013; 19(8): 1253-9.

Rosin P, Niskanen T, Palm D, Struelens M, Takkinen J; Shiga toxin-producing *Escherichia coli* Experts of European Union Food- and Waterborne Diseases and Zoonoses Network. Laboratory preparedness for detection and monitoring of Shiga toxin 2-producing *Escherichia coli* O104:H4 in Europe and response to the 2011 outbreak. *Euro Surveill.* 2013; 18(25).

Scharlach M, Diercke M, Dreesman J, Jahn N, Kriek M, Beyrer K, Claußen K, Pulz M, Floride R. Epidemiological analysis of a cluster within the outbreak of Shiga toxin-producing *Escherichia coli* serotype O104:H4 in Northern Germany, 2011. *Int J Hyg Environ Health.* 2013; 216(3): 341-5.

Soon JM, Seaman P, Baines RN. *Escherichia coli* O104:H4 outbreak from sprouted seeds. *Int J Hyg Environ Health.* 2013; 216(3): 346-54.

Szmolka A, Nagy B. Multidrug resistant commensal *Escherichia coli* in animals and its impact for public health. *Front Microbiol.* 2013; 4: 258. Review.

Taylor EV, Nguyen TA, Machesky KD, Koch E, Sotir MJ, Bohm SR, Folster JP, Bokanyi R, Kupper A, Bidol SA, Emanuel A, Arends KD, Johnson SA, Dunn J, Stroika S, Patel MK, Williams I. Multistate outbreak of *Escherichia coli* O145 infections associated with romaine lettuce consumption, 2010. *J Food Prot.* 2013 Jun; 76(6): 939-44.

Wang J, Stephan R, Karczmarczyk M, Yan Q, Hächler H, Fanning S. Molecular characterization of bla ESBL-harboring conjugative plasmids identified in multi-drug resistant *Escherichia coli* isolated from food-producing animals and healthy humans. *Front Microbiol.* 2013; 4: 188.

Wedley AL, Elajnef HM, Fletcher JN. Characterization of a novel EAST-negative enteropathogenic *E. coli* strain implicated in a food-borne outbreak of diarrhoea in adults. *APMIS.* 2013; 121(6): 494-502.

Won G, Schlegel PJ, Schrock JM, LeJeune JT. Absence of direct association between coliforms and *Escherichia coli* in irrigation water and on produce. *J Food Prot.* 2013; 76(6): 959-66.

Zweifel C, Cernela N, Stephan R. Detection of the Emerging Shiga Toxin-Producing *Escherichia coli* O26:H11/H- Sequence Type 29 (ST29) Clone in Human Patients and Healthy Cattle in Switzerland. *Appl Environ Microbiol.* 2013; 79(17): 5411-3.

Hepatitis A

Lee JH, Lee GC, Kim JI, Yi HA, Lee CH. Development of a new cell culture-based method and optimized protocol for the detection of enteric viruses. *J Virol Methods.* 2013; 191(1): 16-23.

Marrs T, Bruce KD, Logan K, Rivett DW, Perkin MR, Lack G, Flohr C. Is there an association between microbial exposure and food allergy? A systematic review. *Pediatr Allergy Immunol.* 2013; 24(4): 311-320.e8.

Seo JY, Choi BY, Ki M, Jang HL, Park HS, Son HJ, Bae SH, Kang JH, Jun DW, Lee JW, Hong YJ, Kim YS, Kim CH, Chang UI, Kim JH, Yang HW, Kim HS, Park KB, Hwang JS, Heo J, Kim IH, Kim JS, Cheon GJ. Risk factors for acute hepatitis A infection in Korea in 2007 and 2009: a case-control study. *J Korean Med Sci.* 2013; 28(6): 908-14.

Hepatitis E

Abravanel F, Lhomme S, Dubois M, Peron JM, Alric L, Kamar N, Izopet J. Hepatitis E virus. *Med Mal Infect.* 2013; 43(7): 263-70.

Christou L, Kosmidou M. Hepatitis E virus in the Western world--a pork-related zoonosis. *Clin Microbiol Infect.* 2013; 19(7): 600-4.

Cleland A, Smith L, Crossan C, Blatchford O, Dalton HR, Scobie L, Petrik J. Hepatitis E virus in Scottish blood donors. *Vox Sang.* 2013 Jun 13. doi: 10.1111/vox.12056. [Epub ahead of print]

Deshpande A, Gunson R, Bradley A, Cameron S. Hepatitis E: the West of Scotland experience. *Scott Med J.* 2013; 58(3): 178-81.

Gupta E, Pandey P, Pandey S, Sharma MK, Sarin SK. Role of hepatitis E virus antigen in confirming active viral replication in patients with acute viral hepatitis E infection. *J Clin Virol.* 2013 Aug 9. doi:pii: S1386-6532(13)00323-5. 10.1016/j.jcv.2013.07.019. [Epub ahead of print]

Ma H, Geng Y, Li Z, Harrison TJ, Huang W, Zhao C, Wang Y. Analysis of the complete genome sequences of one swine and two human hepatitis E virus genotype 4 strains isolated in Beijing, China. *Infect Genet Evol.* 2013; 18:42-7.

Okano H, Takahashi M, Isono Y, Tanaka H, Nakano T, Oya Y, Sugimoto K, Ito K, Ohmori S, Maegawa T, Kobayashi M, Nagashima S, Nishizawa T, Okamoto H. Characterization of sporadic acute hepatitis E and comparison of hepatitis E virus genomes in acute hepatitis patients and pig liver sold as food in Mie, Japan. *Hepatol Res.* 2013 Aug 8. doi: 10.1111/hepr.12216. [Epub ahead of print]

Pérez-Gracia MT, Lindemann ML, Villalba MC. Hepatitis E: current status. *Rev Med Virol.* 2013 Aug 28. doi: 10.1002/rmv.1759. [Epub ahead of print]

Slot E, Hogema B, Riezebos-Brilman A, Kok T, Molier M, Zaaijer H. Silent hepatitis E virus infection in Dutch blood donors, 2011 to 2012. *Euro Surveill.* 2013; 18(31). doi:pii: 20550.

Wang S, Cheng X, Dai X, Dong C, Xu M, Liang J, Dong M, Purdy MA, Meng J. Rabbit and human hepatitis E virus strains belong to a single serotype. *Virus Res.* 2013; 176(1-2): 101-6.

Listeria monocytogenes

Alali WQ, Schaffner DW. Relationship between *Listeria monocytogenes* and *Listeria* spp. in seafood processing plants. *J Food Prot.* 2013; 76(7): 1279-82.

Ben Slama R, Miladi H, Chaieb K, Bakhrouf A. Survival of *Listeria monocytogenes* cells and the effect of extended frozen storage (-20°C) on the expression of its virulence gene. *Appl Biochem Biotechnol.* 2013; 170(5): 1174-83.

Bodro M, Paterson DL. Listeriosis in patients receiving biologic therapies. *Eur J Clin Microbiol Infect Dis.* 2013; 32(9): 1225-30.

Breidt F Jr, Kay K, Cook J, Osborne J, Ingham B, Arritt F. Determination of 5-log reduction times for *Escherichia coli* O157:H7, *Salmonella enterica*, or *Listeria monocytogenes* in acidified foods with pH 3.5 or 3.8. *J Food Prot.* 2013; 76(7): 1245-9.

Cantinelli T, Chenal-Francisque V, Diancourt L, Frezal L, Leclercq A, Wirth T, Lecuit M, Brisse S. Epidemic clones of *Listeria monocytogenes* are widespread and ancient clonal groups. *J Clin Microbiol.* 2013 Sep 4. [Epub ahead of print] PubMed PMID: 24006010.

Centers for Disease Control and Prevention (CDC). Vital signs: *Listeria* illnesses, deaths, and outbreaks--United States, 2009-2011. *MMWR Morb Mortal Wkly Rep.* 2013; 62(22): 448-52.

Durack J, Ross T, Bowman JP. Characterisation of the Transcriptomes of Genetically Diverse *Listeria monocytogenes* Exposed to Hyperosmotic and Low Temperature Conditions Reveal Global Stress-Adaptation Mechanisms. *PLoS One.* 2013; 8(9):e73603.

Ferrier R, Hezard B, Lintz A, Stahl V, Augustin JC. Combining Individual-Based Modeling and Food Microenvironment Descriptions To Predict the Growth of *Listeria monocytogenes* on Smear Soft Cheese. *Appl Environ Microbiol.* 2013;79(19): 5870-81.

Félix B, Niskanen T, Vingadassalon N, Dao TT, Asséré A, Lombard B, Brisabois A, Roussel S. Pulsed-Field Gel Electrophoresis Proficiency Testing Trials: Toward European Harmonization of the Typing of Food and Clinical Strains of *Listeria monocytogenes*. *Foodborne Pathog Dis.* 2013 Aug 17. [Epub ahead of print] PubMed PMID: 23952473.

Hoelzer K, Chen Y, Dennis S, Evans P, Pouillot R, Silk BJ, Walls I. New Data, Strategies, and Insights for *Listeria monocytogenes* Dose-Response Models: Summary of an Interagency Workshop, 2011. *Risk Anal.* 2013; 33(9): 1568-81.

Hystead E, Diez-Gonzalez F, Schoenfuss TC. The effect of sodium reduction with and without potassium chloride on the survival of *Listeria monocytogenes* in Cheddar cheese. *J Dairy Sci.* 2013 Aug 1. doi:pil: S0022-0302(13)00529-8. 10.3168/jds.2013-6675. [Epub ahead of print]

Kim MJ, Kim SA, Kang YS, Hwang IG, Rhee MS. Microbial diversity and prevalence of foodborne pathogens in cheap and junk foods consumed by primary schoolchildren. *Lett Appl Microbiol.* 2013; 57(1): 47-53.

Lamden KH, Fox AJ, Amar CF, Little CL. A case of foodborne listeriosis linked to a contaminated food production process. *J Med Microbiol.* 2013 Jun 20. [Epub ahead of print] PubMed PMID: 23788596.

Lazarus C, Leclercq A, Lecuit M, Vaillant V, Coignard B, Blanchard H, Novakova I, Astagneau P. Probable nosocomial transmission of listeriosis in neonates. *J Hosp Infect.* 2013 Aug 31. doi:pil: S0195-6701(13)00261-2. 10.1016/j.jhin.2013.06.019. [Epub ahead of print]

Lobacz A, Kowalik J, Tarczyska A. Modeling the growth of *Listeria monocytogenes* in mold-ripened cheeses. *J Dairy Sci.* 2013; 96(6): 3449-60.

McCollum JT, Cronquist AB, Silk BJ, Jackson KA, O'Connor KA, Cosgrove S, Gossack JP, Parachini SS, Jain NS, Ettestad P, Ibraheem M, Cantu V, Joshi M, DuVernoy T, Fogg NW Jr, Gorny JR, Mogen KM, Spires C, Teitell P, Joseph LA, Tarr CL, Imanishi M, Neil KP, Tauxe RV, Mahon BE. Multistate outbreak of listeriosis associated with cantaloupe. *N Engl J Med.* 2013 Sep 5;369(10):944-53.

Meldrum RJ, Charles D, Mannion P, Ellis P. Variation in the microbiological quality of commercially produced vacuum-packed cooked sliced meat between production and the end of shelf-life. *Int J Environ Health Res.* 2013 Jul 22. [Epub ahead of print] PubMed PMID: 23869397.

Montiel R, Bravo D, Medina M. Commercial Biopreservatives Combined with Salt and Sugar To Control *Listeria monocytogenes* during Smoked Salmon Processing. *J Food Prot.* 2013; 76(8): 1463-5.

Poulsen KP, Czuprynski CJ. Pathogenesis of listeriosis during pregnancy. *Anim Health Res Rev.* 2013; 14(1): 30-9.

Saleh-Lakha S, Allen VG, Li J, Pagotto F, Odumeru J, Taboada E, Lombos M, Tabing KC, Blais B, Ogunremi D, Downing G, Lee S, Gao A, Nadon C, Chen S. Subtyping of a large collection of historical *Listeria monocytogenes* strains from Ontario, Canada, using an improved Multiple-Locus Variable-Number Tandem Repeat Analysis (MLVA). *Appl Environ Microbiol*. 2013 Aug 16. [Epub ahead of print] PubMed PMID: 23956391.

Sansawat T, Zhang L, Jeong JY, Xu Y, Hessel GW, Ryser ET, Harte JB, Tempelman R, Kang I. Inhibition of *Listeria monocytogenes* in Full- and Low-Sodium Frankfurters at 4, 7, or 10°C. Using Spray-Dried Mixtures of Organic Acid Salts. *J Food Prot*. 2013; 76(9): 1557-67.

Sasaki Y, Haruna M, Murakami M, Hayashida M, Takahashi N, Urushiyama T, Ito K, Yamada Y. Contamination of Poultry Products with *Listeria monocytogenes* at Poultry Processing Plants. *J Vet Med Sci*. 2013 Sep 9. [Epub ahead of print] PubMed PMID: 24018827.

Schaffner DW. Utilization of mathematical models to manage risk of holding cold food without temperature control. *J Food Prot*. 2013; 76(6): 1085-94.

Shen J, Rump L, Zhang Y, Chen Y, Wang X, Meng J. Molecular subtyping and virulence gene analysis of *Listeria monocytogenes* isolates from food. *Food Microbiol*. 2013; 35(1): 58-64.

Tenenhaus-Aziza F, Daudin JJ, Maffre A, Sanaa M. Risk-Based Approach for Microbiological Food Safety Management in the Dairy Industry: The Case of *Listeria monocytogenes* in Soft Cheese Made from Pasteurized Milk. *Risk Anal*. 2013 Jun 18. doi: 10.1111/risa.12074. [Epub ahead of print]

Norovirus

Bitler EJ, Matthews JE, Dickey BW, Eisenberg JN, Leon JS. Norovirus outbreaks: a systematic review of commonly implicated transmission routes and vehicles. *Epidemiol Infect*. 2013;141(8):1563-71.

Dicaprio E, Ma Y, Hughes J, Li J. Epidemiology, prevention, and control of the number one foodborne illness: human norovirus. *Infect Dis Clin North Am*. 2013; 27(3): 651-74.

El-Senousy WM, Costafreda MI, Pintó RM, Bosch A. Method validation for norovirus detection in naturally contaminated irrigation water and fresh produce. *Int J Food Microbiol*. 2013 Jul 1. doi:pii: S0168-1605(13)00314-0. 10.1016/j.ijfoodmicro.2013.06.023. [Epub ahead of print]

- Gaulin C, Nguon S, Leblanc MA, Ramsay D, Roy S. Multiple outbreaks of gastroenteritis that were associated with 16 funerals and a unique caterer and spanned 6 days, 2011, Québec, Canada. *J Food Prot.* 2013; 76(9): 1582-9.
- Hoelzer K, Fanaselle W, Pouillot R, Van Doren JM, Dennis S. Virus inactivation on hard surfaces or in suspension by chemical disinfectants: systematic review and meta-analysis of norovirus surrogates. *J Food Prot.* 2013; 76(6): 1006-16.
- Hoffmann D, Mauroy A, Seebach J, Simon V, Wantia N, Protzer U. New norovirus classified as a recombinant GII.g/GII.1 causes an extended foodborne outbreak at a university hospital in Munich. *J Clin Virol.* 2013; 58(1): 24-30.
- Lee BR, Lee SG, Park JH, Kim KY, Ryu SR, Rhee OJ, Park JW, Lee JS, Paik SY. Norovirus contamination levels in ground water treatment systems used for food-catering facilities in South Korea. *Viruses.* 2013; 5(7): 1646-54.
- Manso CF, Romalde JL. Detection and characterization of hepatitis A virus and norovirus in mussels from Galicia (NW Spain). *Food Environ Virol.* 2013; 5(2): 110-8.
- Maritschnik S, Kanitz EE, Simons E, Höhne M, Neumann H, Allerberger F, Schmid D, Lederer I. A Food Handler-Associated, Foodborne Norovirus GII.4 Sydney 2012-Outbreak Following a Wedding Dinner, Austria, October 2012. *Food Environ Virol.* 2013 Sep 12. [Epub ahead of print] PubMed PMID: 24026524.
- Mäde D, Trübner K, Neubert E, Höhne M, Johne R. Detection and Typing of Norovirus from Frozen Strawberries Involved in a Large-Scale Gastroenteritis Outbreak in Germany. *Food Environ Virol.* 2013 Jul 26. [Epub ahead of print] PubMed PMID: 23888384.
- Pavoni E, Consoli M, Suffredini E, Arcangeli G, Serracca L, Battistini R, Rossini I, Croci L, Losio MN. Noroviruses in seafood: a 9-year monitoring in Italy. *Foodborne Pathog Dis.* 2013; 10(6): 533-9.
- Repp KK, Hostetler TP, Keene WE. A norovirus outbreak related to contaminated surfaces. *J Infect Dis.* 2013; 208(2): 295-8.
- Ruan F, Tan AJ, Man TF, Li H, Mo YL, Lin YX, Deng XL. Gastroenteritis Outbreaks Caused by Norovirus Genotype II.7 in a College in China (Zhuhai, Guangdong) in 2011. *Foodborne Pathog Dis.* 2013 Jul 10. [Epub ahead of print] PubMed PMID: 23841656.
- Rönnqvist M, Rättö M, Tuominen P, Salo S, Maunula L. Swabs as a tool for monitoring the presence of norovirus on environmental surfaces in the food industry. *J Food Prot.* 2013; 76(8): 1421-8.

Schaeffer J, Le Saux JC, Lora M, Atmar RL, Le Guyader FS. Norovirus contamination on French marketed oysters. *Int J Food Microbiol.* 2013; 166(2):244-8.

Stals A, Uyttendaele M, Baert L, Van Coillie E. Norovirus transfer between foods and food contact materials. *J Food Prot.* 2013; 76(7): 1202-9.

Thornley CN, Hewitt J, Perumal L, Van Gessel SM, Wong J, David SA, Rapana JP, Li S, Marshall JC, Greening GE. Multiple outbreaks of a novel norovirus GII.4 linked to an infected post-symptomatic food handler. *Epidemiol Infect.* 2013; 141(8):1585-97.

Tung G, Macinga D, Arbogast J, Jaykus LA. Efficacy of commonly used disinfectants for inactivation of human noroviruses and their surrogates. *J Food Prot.* 2013; 76(7):1210-7.

Verhaelen K, Bouwknecht M, Carratalà A, Lodder-Verschoor F, Diez-Valcarce M, Rodríguez-Lázaro D, de Roda Husman AM, Rutjes SA. Virus transfer proportions between gloved fingertips, soft berries, and lettuce, and associated health risks. *Int J Food Microbiol.* 2013; 166(3): 419-425.

Wang Q, Hirneisen K, Markland S, Kniel KE. Survival of murine norovirus, Tulane virus and Hepatitis A virus on alfalfa seeds and sprouts during storage and germination. *Appl Environ Microbiol.* 2013 Sep 6. [Epub ahead of print]

Salmonella

Bassal R, Reisfeld A, Nissan I, Agmon V, Taran D, Schemberg B, Cohen D, Shohat T. Risk factors for sporadic infection with *Salmonella Infantis*: a matched case-control study. *Epidemiol Infect.* 2013 Jul 23:1-6. [Epub ahead of print] PubMed PMID: 23879895.

Batz MB, Henke E, Kowalcyk B. Long-term consequences of foodborne infections. *Infect Dis Clin North Am.* 2013; 27(3): 599-616.

Berriman AD, Clancy D, Clough HE, Armstrong D, Christley RM. Effectiveness of simulated interventions in reducing the estimated prevalence of *Salmonella* in UK pig herds. *PLoS One.* 2013; 8(6):e66054.

Bouguelia S, Roupioz Y, Slimani S, Mondani L, Casabona MG, Durmort C, Vernet T, Calemczuk R, Livache T. On-chip microbial culture for the specific detection of very low levels of bacteria. *Lab Chip.* 2013 Aug 2. [Epub ahead of print] PubMed PMID: 23912527.

Bouwknegt M, van Pelt W, Havelaar AH. Scoping the impact of changes in population age-structure on the future burden of foodborne disease in the Netherlands, 2020-2060. *Int J Environ Res Public Health*. 2013; 10(7): 2888-96.

Breidt F Jr, Kay K, Cook J, Osborne J, Ingham B, Arritt F. Determination of 5-log reduction times for *Escherichia coli* O157:H7, *Salmonella enterica*, or *Listeria monocytogenes* in acidified foods with pH 3.5 or 3.8. *J Food Prot*. 2013; 76(7): 1245-9.

Campos J, Mourão J, Pestana N, Peixe L, Novais C, Antunes P. Microbiological quality of ready-to-eat salads: An underestimated vehicle of bacteria and clinically relevant antibiotic resistance genes. *Int J Food Microbiol*. 2013; 166(3): 464-470.

Centers for Disease Control and Prevention (CDC). Outbreak of *Salmonella* Heidelberg infections linked to a single poultry producer -- 13 states, 2012-2013. *MMWR Morb Mortal Wkly Rep*. 2013; 62(27): 553-6.

Chai Y, Horikawa S, Li S, Wikle HC, Chin BA. A surface-scanning coil detector for real-time, in-situ detection of bacteria on fresh food surfaces. *Biosens Bioelectron*. 2013; 50: 311-7.

Dahl J. Controlling *Salmonella* in live pigs and at slaughter: The Danish experience. *Vet J*. 2013 Sep 12. doi:pii: S1090-0233(13)00244-X. 10.1016/j.tvjl.2013.05.028. [Epub ahead of print]

Dey M, Mayo JA, Saville D, Wolyniak C, Klontz KC. Recalls of foods due to microbiological contamination classified by the U.S. Food and Drug Administration, fiscal years 2003 through 2011. *J Food Prot*. 2013; 76(6): 932-8.

Economou V, Gousia P, Kansouzidou A, Sakkas H, Karanis P, Papadopoulou C. Prevalence, antimicrobial resistance and relation to indicator and pathogenic microorganisms of *Salmonella enterica* isolated from surface waters within an agricultural landscape. *Int J Hyg Environ Health*. 2013; 216(4): 435-44.

Elkins CA, Kotewicz ML, Jackson SA, Lacher DW, Abu-Ali GS, Patel IR. Genomic paradigms for food-borne enteric pathogen analysis at the USFDA: case studies highlighting method utility, integration and resolution. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess*. 2013; 30(8): 1422-36.

Fernandes RD, Freire MT, de Paula ES, Kanashiro AL, Catunda FA, Rosa AF, Balieiro JC, Trindade MA. Stability of lamb loin stored under refrigeration and packed in different modified atmosphere packaging systems. *Meat Sci*. 2013; 96(1):554-561.

Finn S, Hinton JC, McClure P, Amézquita A, Martins M, Fanning S. Phenotypic

characterization of salmonella isolated from food production environments associated with low-water activity foods. *J Food Prot.* 2013; 76(9): 1488-99.

Finn S, Händler K, Condell O, Colgan A, Cooney S, McClure P, Amézquita A, Hinton JC, Fanning S. ProP is required for the survival of desiccated *Salmonella enterica* serovar typhimurium cells on a stainless steel surface. *Appl Environ Microbiol.* 2013; 79(14): 4376-84.

Ge B, Lafon PC, Carter PJ, McDermott SD, Abbott J, Glenn A, Ayers SL, Friedman SL, Paige JC, Wagner DD, Zhao S, McDermott PF, Rasmussen MA. Retrospective Analysis of *Salmonella*, *Campylobacter*, *Escherichia coli*, and *Enterococcus* in Animal Feed Ingredients. *Foodborne Pathog Dis.* 2013; 10(8): 684-91.

Gieraltowski L, Julian E, Pringle J, Macdonald K, Quilliam D, Marsden-Haug N, Saathoff-Huber L, Von Stein D, Kissler B, Parish M, Elder D, Howard-King V, Besser J, Sodha S, Loharikar A, Dalton S, Williams I, Barton Behravesh C. Nationwide outbreak of *Salmonella* Montevideo infections associated with contaminated imported black and red pepper: warehouse membership cards provide critical clues to identify the source. *Epidemiol Infect.* 2013; 141(6): 1244-52.

Giombelli A, Cavani R, Gloria MB. Evaluation of three sampling methods for the microbiological analysis of broiler carcasses after immersion chilling. *J Food Prot.* 2013; 76(8): 1330-5.

Glenn LM, Lindsey RL, Folster JP, Pecic G, Boerlin P, Gilmour MW, Harbottle H, Zhao S, McDermott PF, Fedorka-Cray PJ, Frye JG. Antimicrobial resistance genes in multidrug-resistant *Salmonella enterica* isolated from animals, retail meats, and humans in the United States and Canada. *Microb Drug Resist.* 2013; 19(3): 175-84.

Gopinath G, Hari K, Jain R, Mammel MK, Kothary MH, Franco AA, Grim CJ, Jarvis KG, Sathyamoorthy V, Hu L, Datta AR, Patel IR, Jackson SA, Gangiredla J, Kotewicz ML, LeClerc JE, Wekell M, McCardell BA, Solomotis MD, Tall BD. The Pathogen-annotated Tracking Resource Network (PATRN) system: a web-based resource to aid food safety, regulatory science, and investigations of foodborne pathogens and disease. *Food Microbiol.* 2013; 34(2): 303-18.

Gould LH, Walsh KA, Vieira AR, Herman K, Williams IT, Hall AJ, Cole D; Centers for Disease Control and Prevention. Surveillance for foodborne disease outbreaks - United States, 1998-2008. *MMWR Surveill Summ.* 2013; 62(2): 1-34.

Guillier L, Danan C, Bergis H, Delignette-Muller ML, Granier S, Rudelle S, Beaufort A, Brisabois A. Use of quantitative microbial risk assessment when investigating foodborne illness outbreaks: The example of a monophasic *Salmonella*

Typhimurium 4,5,12:i:- outbreak implicating beef burgers. *Int J Food Microbiol.* 2013; 166(3): 471-478.

Hara-Kudo Y, Konuma H, Kamata Y, Miyahara M, Takatori K, Onoue Y, Sugita-Konishi Y, Ohnishi T. Prevalence of the main food-borne pathogens in retail food under the national food surveillance system in Japan. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess.* 2013; 30(8): 1450-8.

Hardy B, Crilly N, Pendleton S, Andino A, Wallis A, Zhang N, Hanning I. Impact of rearing conditions on the microbiological quality of raw retail poultry meat. *J Food Sci.* 2013; 78(8):M1232-5.

He Y, Li Y, Salazar JK, Yang J, Tortorello ML, Zhang W. Increased water activity reduces the thermal resistance of *Salmonella enterica* in peanut butter. *Appl Environ Microbiol.* 2013; 79(15): 4763-7.

Jackson BR, Griffin PM, Cole D, Walsh KA, Chai SJ. Outbreak-associated *Salmonella enterica* serotypes and food Commodities, United States, 1998-2008. *Emerg Infect Dis.* 2013; 19(8): 1239-44.

Jensen DA, Friedrich LM, Harris LJ, Danyluk MD, Schaffner DW. Quantifying Transfer Rates of *Salmonella* and *Escherichia coli* O157:H7 between Fresh-Cut Produce and Common Kitchen Surfaces. *J Food Prot.* 2013; 76(9): 1530-8.

Kostoulas P, Nielsen SS, Browne WJ, Leontides L. Sample size estimation to substantiate freedom from disease for clustered binary data with a specific risk profile. *Epidemiol Infect.* 2013; 141(6): 1318-27.

Le Hello S, Harrois D, Bouchrif B, Sontag L, Elhani D, Guibert V, Zerouali K, Weill FX. Highly drug-resistant *Salmonella enterica* serotype Kentucky ST198-X1: a microbiological study. *Lancet Infect Dis.* 2013; 13(8): 672-9.

Li D, Friedrich LM, Danyluk MD, Harris LJ, Schaffner DW. Development and validation of a mathematical model for growth of pathogens in cut melons. *J Food Prot.* 2013; 76(6): 953-8.

Lianou A, Koutsoumanis KP. Evaluation of the strain variability of *Salmonella enterica* acid and heat resistance. *Food Microbiol.* 2013; 34(2): 259-67.

Martins M, McCusker MP, McCabe EM, O'Leary D, Duffy G, Fanning S. Evidence of Metabolic Switching and Implications for Food Safety from the Phenome(s) of *Salmonella enterica* Serovar Typhimurium DT104 Cultured at Selected Points across the Pork Production Food Chain. *Appl Environ Microbiol.* 2013; 79(18): 5437-49.

Menconi A, Shivaramaiah S, Huff GR, Prado O, Morales JE, Pumford NR, Morgan M, Wolfenden A, Bielke LR, Hargis BM, Tellez G. Effect of different concentrations of acetic, citric, and propionic acid dipping solutions on bacterial contamination of raw chicken skin. *Poult Sci.* 2013; 92(8): 2216-20.

Mughini-Gras L, Barrucci F, Smid JH, Graziani C, Luzzi I, Ricci A, Barco L, Rosmini R, Havelaar AH, VAN Pelt W, Busani L. Attribution of human *Salmonella* infections to animal and food sources in Italy (2002-2010): adaptations of the Dutch and modified Hald source attribution models. *Epidemiol Infect.* 2013 Aug 7:1-13. [Epub ahead of print]

Møller CO, Ilg Y, Aabo S, Christensen BB, Dalgaard P, Hansen TB. Effect of natural microbiota on growth of *Salmonella* spp. in fresh pork--a predictive microbiology approach. *Food Microbiol.* 2013; 34(2): 284-95.

Nagel GM, Bauermeister LJ, Bratcher CL, Singh M, McKee SR. *Salmonella* and *Campylobacter* reduction and quality characteristics of poultry carcasses treated with various antimicrobials in a post-chill immersion tank. *Int J Food Microbiol.* 2013; 165(3): 281-6.

Nauta M, Barfod K, Hald T, Sørensen AH, Emborg HD, Aabo S. Prediction of *Salmonella* carcass contamination by a comparative quantitative analysis of *E. coli* and *Salmonella* during pig slaughter. *Int J Food Microbiol.* 2013; 166(2): 231-7.

Parmley EJ, Pintar K, Majowicz S, Avery B, Cook A, Jokinen C, Gannon V, Lapen DR, Topp E, Edge TA, Gilmour M, Pollari F, Reid-Smith R, Irwin R. A canadian application of one health: integration of salmonella data from various canadian surveillance programs (2005-2010). *Foodborne Pathog Dis.* 2013; 10(9): 747-56.

Quigley L, O'Sullivan O, Stanton C, Beresford TP, Ross RP, Fitzgerald GF, Cotter PD. The complex microbiota of raw milk. *FEMS Microbiol Rev.* 2013; 37(5): 664-98.

Quinlan JJ. Foodborne illness incidence rates and food safety risks for populations of low socioeconomic status and minority race/ethnicity: a review of the literature. *Int J Environ Res Public Health.* 2013; 10(8): 3634-52.

Racicot M, Kocher A, Beauchamp G, Letellier A, Vaillancourt JP. Assessing most practical and effective protocols to sanitize hands of poultry catching crew members. *Prev Vet Med.* 2013; 111(1-2): 92-9.

Rathgeber BM, McCarron P, Budgell KL. *Salmonella* penetration through eggshells of chickens of different genetic backgrounds. *Poult Sci.* 2013; 92(9): 2457-62.

Rhoades J, Kargiotou C, Katsanidis E, Koutsoumanis KP. Use of marination for

controlling *Salmonella enterica* and *Listeria monocytogenes* in raw beef. *Food Microbiol.* 2013; 36(2): 248-53.

Ricke SC, Khatiwara A, Kwon YM. Application of microarray analysis of foodborne *Salmonella* in poultry production: a review. *Poult Sci.* 2013; 92(9): 2243-50.

Santillana Farakos SM, Frank JF, Schaffner DW. Modeling the influence of temperature, water activity and water mobility on the persistence of *Salmonella* in low-moisture foods. *Int J Food Microbiol.* 2013; 166(2): 280-93.

Schaffner DW. Utilization of mathematical models to manage risk of holding cold food without temperature control. *J Food Prot.* 2013; 76(6): 1085-94.

Schelin J, Andersson G, Vigre H, Norling B, Häggblom P, Hoorfar J, Rådström P, Löfström C. Evaluation of pre-PCR processing approaches for enumeration of *Salmonella enterica* in naturally contaminated animal feed. *J Appl Microbiol.* 2013 Sep 3. doi: 10.1111/jam.12337. [Epub ahead of print]

Sjölund-Karlsson M, Howie RL, Blickenstaff K, Boerlin P, Ball T, Chalmers G, Duval B, Haro J, Rickert R, Zhao S, Fedorka-Cray PJ, Whichard JM. Occurrence of β -lactamase genes among non-Typhi *Salmonella enterica* isolated from humans, food animals, and retail meats in the United States and Canada. *Microb Drug Resist.* 2013; 19(3): 191-7.

Spricigo DA, Bardina C, Cortés P, Llagostera M. Use of a bacteriophage cocktail to control *Salmonella* in food and the food industry. *Int J Food Microbiol.* 2013; 165(2): 169-74.

Thomas MK, Murray R, Flockhart L, Pintar K, Pollari F, Fazil A, Nesbitt A, Marshall B. Estimates of the burden of foodborne illness in Canada for 30 specified pathogens and unspecified agents, circa 2006. *Foodborne Pathog Dis.* 2013; 10(7): 639-48.

Tian JQ, Bae YM, Lee SY. Survival of foodborne pathogens at different relative humidities and temperatures and the effect of sanitizers on apples with different surface conditions. *Food Microbiol.* 2013; 35(1): 21-6.

Zulfakar SS, White JD, Ross T, Tamplin M. Effect of pH, salt and chemical rinses on bacterial attachment to extracellular matrix proteins. *Food Microbiol.* 2013; 34(2): 369-75.

Staphylococcus aureus

Ebner R, Johler S, Sihto HM, Stephan R, Zweifel C. Microarray-Based Characterization of *Staphylococcus aureus* Isolates Obtained from Chicken Carcasses. *J Food Prot.* 2013; 76(8): 1471-4.

Ho J, O'Donoghue MM, Boost MV. Occupational exposure to raw meat: A newly-recognized risk factor for *Staphylococcus aureus* nasal colonization amongst food handlers. *Int J Hyg Environ Health.* 2013 Jul 24. doi:pii: S1438-4639(13)00103-X. 10.1016/j.ijheh.2013.07.009. [Epub ahead of print]

Lassok B, Tenhagen BA. From pig to pork: methicillin-resistant *Staphylococcus aureus* in the pork production chain. *J Food Prot.* 2013; 76(6):1095-108.

Martins PD, de Almeida TT, Basso AP, de Moura TM, Frazzon J, Tondo EC, Frazzon AP. Coagulase-positive *staphylococci* isolated from chicken meat: pathogenic potential and vancomycin resistance. *Foodborne Pathog Dis.* 2013; 10(9): 771-6.

Mole B. MRSA: Farming up trouble. *Nature.* 2013; 499(7459):398-400.

Nemeghaire S, Roelandt S, Argudín MA, Haesebrouck F, Butaye P. Characterization of methicillin-resistant *Staphylococcus aureus* from healthy carrier chickens. *Avian Pathol.* 2013; 42(4): 342-6.

Verheghe M, Pletinckx LJ, Crombé F, Vandersmissen T, Haesebrouck F, Butaye P, Heyndrickx M, Rasschaert G. Methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 in pig farms and multispecies farms. *Zoonoses Public Health.* 2013; 60(5): 366-74.

Toxoplasma gondii

Bellali H, Pelloux H, Villena I, Fricker-Hidalgo H, Le Strat Y, Goulet V. Prevalence of toxoplasmosis in France in 1998: Is there a difference between men and women? At what age do children become infected? *Rev Epidemiol Sante Publique.* 2013; 61(4): 311-7.

Blackman MJ, Carruthers VB. Recent insights into apicomplexan parasite egress provide new views to a kill. *Curr Opin Microbiol.* 2013; 16(4): 459-64.

Burrells A, Bartley PM, Zimmer IA, Roy S, Kitchener AC, Meredith A, Wright SE, Innes EA, Katzer F. Evidence of the three main clonal *Toxoplasma gondii* lineages from wild mammalian carnivores in the UK. *Parasitology.* 2013 Aug 19:1-9. [Epub ahead of print].

Chumpolbanchorn K, Lymbery AJ, Pallant LJ, Pan S, Sukthana Y, Thompson RC. A high prevalence of *Toxoplasma* in Australian chickens. *Vet Parasitol.* 2013; 196(1-2): 209-11.

Djurković-Djaković O, Bobić B, Nikolić A, Klun I, Dupouy-Camet J. Pork as a source of human parasitic infection. *Clin Microbiol Infect.* 2013; 19(7): 586-94.

Flatt A, Shetty N. Seroprevalence and risk factors for toxoplasmosis among antenatal women in London: a re-examination of risk in an ethnically diverse population. *Eur J Public Health.* 2013; 23(4): 648-52.

García-Bocanegra I, Cabezón O, Hernández E, Martínez-Cruz MS, Martínez-Moreno Á, Martínez-Moreno J. *Toxoplasma gondii* in ruminant species (cattle, sheep, and goats) from southern Spain. *J Parasitol.* 2013; 99(3): 438-40.

Hamdani N, Daban-Huard C, Lajnef M, Richard JR, Delavest M, Godin O, Le Guen E, Vederine FE, Lépine JP, Jamain S, Houenou J, Le Corvoisier P, Aoki M, Moins-Teisserenc H, Charron D, Krishnamoorthy R, Yolken R, Dickerson F, Tamouza R, Leboyer M. Relationship between *Toxoplasma gondii* infection and bipolar disorder in a French sample. *J Affect Disord.* 2013; 148(2-3): 444-8.

Kemp LE, Yamamoto M, Soldati-Favre D. Subversion of host cellular functions by the apicomplexan parasites. *FEMS Microbiol Rev.* 2013; 37(4): 607-31.

Lélu M, Langlais M, Poulle ML, Gilot-Fromont E, Gandon S. When should a trophically and vertically transmitted parasite manipulate its intermediate host? The case of *Toxoplasma gondii*. *Proc Biol Sci.* 2013; 280(1765): 20131143.

Pott S, Koethe M, Bangoura B, Zöller B, Dauschies A, Straubinger RK, Fehlhaber K, Ludewig M. Effects of pH, sodium chloride, and curing salt on the infectivity of *Toxoplasma gondii* tissue cysts. *J Food Prot.* 2013; 76(6): 1056-61.

Rommereim LM, Hortua Triana MA, Falla A, Sanders KL, Guevara RB, Bzik DJ, Fox BA. Genetic Manipulation in $\Delta ku80$ Strains for Functional Genomic Analysis of *Toxoplasma gondii*. *J Vis Exp.* 2013 Jul 12;(77). doi: 10.3791/50598.

Sharma P, Chitnis CE. Key molecular events during host cell invasion by Apicomplexan pathogens. *Curr Opin Microbiol.* 2013; 16(4): 432-7.

Shobab L, Pleyer U, Johnsen J, Metzner S, James ER, Torun N, Fay MP, Liesenfeld O, Grigg ME. *Toxoplasma* Serotype Is Associated With Development of Ocular Toxoplasmosis. *J Infect Dis.* 2013 Aug 22. [Epub ahead of print] PubMed

PMID: 23878321.

Smith G. Food- and water-borne disease: using case control studies to estimate the force of infection that accounts for primary, sporadic cases. *Epidemics*. 2013; 5(2): 77-84.

Torgerson PR, Mastroiacovo P. The global burden of congenital toxoplasmosis: a systematic review. *Bull World Health Organ*. 2013; 91(7): 501-8.

Torrey EF, Yolken RH. *Toxoplasma* oocysts as a public health problem. *Trends Parasitol*. 2013; 29(8): 380-4.

Turner M, Lenhart S, Rosenthal B, Zhao X. Modeling effective transmission pathways and control of the world's most successful parasite. *Theor Popul Biol*. 2013; 86: 50-61.

Turčeková L, Antolová D, Reiterová K, Spišák F. Occurrence and genetic characterization of *Toxoplasma gondii* in naturally infected pigs. *Acta Parasitol*. 2013; 58(3): 361-6.